

NATIONAL ESSENTIAL MEDICINES LIST 2018



**Drug Regulatory Authority of Pakistan
Ministry of National Health Services,
Regulations and Coordination
Government of Pakistan**



Table of Contents

Acronyms.....	v
National Commitment on Essential Medicines	1
Preface	3
Executive Summary	5
Alphabetical List NEML 2018.....	9
National Essential Medicine List of Pakistan 2018.....	17
Medicines with Age and Weight Restriction	57
Index	59
Application Procedure for Inclusion, Deletion or Modification of a Medicine in the Next Edition of National Essential Medicine List, 2020.....	69
References	71

Figures:

Figure 1: Drug Selection	5
Figure 2: WHO Access to Medicines Framework	6

Table:

Table 1: Medicines with Age and Weight Restriction.....	57
---	----

Acronyms

3TC	lamivudine
ABC	abacavir
API	active pharmaceutical ingredient
ATRA	all-trans retinoid acid
D4T	stavudine
DRAP	Drug Regulatory Authority of Pakistan
EML	Essential Medicines List
FDC	fixed-dose combination
GHSC-PSM	Global Health Supply Chain Management Program – Procurement and Supply Management Project
MNCH	maternal, neonatal, and child health
NEML	National Essential Medicines List
NTP	National TB Control Program
P	primary health care level
S	secondary health care level
T	tertiary health care level
TDF	tenofovir disoproxil fumarate
USAID	United States Agency for International Development
U.S.P.	United States Pharmacopeia
WHO	World Health Organization

Message from Federal Minister for National Health Services, Regulations & Coordination

National Essential Medicine List (NEML) is an important element of National Health system policies and practices and has proven to be a useful tool to help country in selecting quality, safe, efficacious and affordable medicines. Availability of NEML is an important task and quality assurance systems benchmark at the national level. The NEML is a critical document that provides a list of essential medicines for primary secondary and tertiary level health facilities. The NEML is based on disease prevalence, evidence of efficacy and safety, comparative cost-effectiveness and parameters adapted by WHO in its latest edition of Model List of Essential Medicines.



Pakistan is committed to aligning its domestic laws and policies with its obligations enshrined in the international treaties and is aspirant to ensure compliance with international best practices. Access to Essential Medicines is part of right to health and important tool to provide access of quality medicines to the people of Pakistan. Cost effectiveness is an important concern for the poor masses. The NEML at hand is also envisaged to take stock of this aspect as well.

I congratulate Drug Regulatory Authority of Pakistan (DRAP), USAID Pakistan, World Health Organization (WHO) and health experts who managed to develop this national document of public importance.

The Government of Pakistan is committed to ensure equity and quality through the delivery of essential preventative and curative care services to every citizen of Pakistan, aligned with the National Health Vision 2016 - 2025.

A handwritten signature in green ink, appearing to read "Aamer Mehmood Kiani".

Mr. Aamer Mehmood Kiani

Federal Minister

National Health Services, Regulations and Coordination, Islamabad

National Commitment on Essential Medicines

October 2018

Essential Medicines are an indispensable component of healthcare system of a country. Pakistan is committed that its domestic law and policies are consistent with its obligations as per international treaties.



Access to essential medicines is a human right. A sizeable population in the developing world lacks access to essential medicines. The availability of affordable and effective essential drugs is, therefore, one of the most visible indicators of the quality of health services. Drug shortages and quality issues continue to undermine the performance of health systems throughout the developing world. Development and implementation of Essential Drugs list plays a vital role in ensuring access of essential medicines to public. We are determined to put our all efforts to ensure the availability of essential medicines.

Development of National Essential Medicines List 2018 is an ongoing tool to provide guidance in the selection of medicines satisfying the criteria of access to safe, quality, and affordable medicines for the people of Pakistan.

Capt Zahid Saeed (Retd)

Secretary

Ministry of National Health Services, Regulations & Coordination

Government of Pakistan

Preface

Under the 18th Constitutional Amendment, health services delivery has been transferred to provinces, whereas Pakistan Vision 2016 – 2025 provides an overarching roadmap for the country which primarily includes reducing the widespread prevalence of communicable diseases, addressing inadequacies in primary, secondary healthcare facilities, improving the pharmaceutical sector to ensure the availability, affordability and quality of drugs and to promote rational use of drugs in the country.



Among the fundamental human rights, vision of essential medicines is one of essential building blocks of health systems. The list of essential medicines is aimed at satisfying the primary healthcare needs of the population. These are intended to be available within the context of functioning of health system at all times, in adequate amounts, in the appropriate dosage form with assured quality, and at a price that the individuals and community can afford.

Careful selection of a limited range of essential medicines of high quality can provide better management towards judicious use of health resources.

National standard treatment guidelines navigate selection of essential medicine with the expectation to ensure their availability with reference to proper use of medicine within the healthcare system while reducing its cost.

National Essential Medicine List NEML, 2018 is a critical document which provides and supports quality of health services across the country.

Department of Pharmacy services, Drug Regulatory Authority of Pakistan was assigned with the responsibility of revising & updating National Essential Medicine List. The Drug Regulatory Authority of Pakistan, Ministry of National Health Services, Regulations & Coordination was assisted by USAID to finalize this National Essential Medicine List for the primary, secondary and tertiary level health facilities. An essential medicines list thus developed is based on disease prevalence, evidence on efficacy and safety, comparative cost-effectiveness and parameters adapted by W.H.O in its latest edition

of Model List of Essential Medicines. WHO's criteria for inclusion and exclusion of medicines was adapted for the purpose. The NEML 2018 includes the medicines needed for adequately addressing the priority health problems as identified in National & Provincial Health strategies.

Health sector in general and Pharmaceutical sector in particular is expected to seriously consider adopting this list as a way forward to guide government in Medicine & Pharmaceutical sector policies, the determinants of medicines access and availability. Pharmacoeconomics parameters in drug selection will guide in drafting health reimbursement strategy and devising reimbursement lists. The Provincial Governments can play a pivotal role to encourage procurement and supply chain management policies based on essential medicine concepts to promote rational use of scarce public resources. Drug Regulatory authority will make all efforts to improve patient health through availability of cost efficacious, safe & quality medicines in country.

The Drug Regulatory Authority of Pakistan, acknowledges and appreciates the contribution of Mr. Sheikh Ansar Ahmad, Director Pharmacy Services Division, DRAP, Prof. Dr. Javed Akram, Ex – Vice Chancellor, Shaheed Zulfiqar Ali Bhutto Medical University, Islamabad and Dr. Muhammad Tariq, Country Director, USAID Global Health Supply Chain Program – Procurement and Supply Management (GHSC-PSM) Project, being implemented by Chemonics International Inc. for his tangible support in upgrading/review and development of NEML 2018 for ensuring governance and accountability within the health system. The Drug Regulatory Authority of Pakistan also acknowledges the technical core committee on NEML, specially the efforts of the Pharmacy Services Division for their human resource and technical input. National list will increase transparency and best value for money in procurement thus leading to increased medicine access and availability in Pakistan.



Dr. Sheikh Akhter Hussain,
Chief Executive Officer,
Drug Regulatory Authority of Pakistan,
Ministry of National Health Services, Regulations and Coordination,
Government of Pakistan.

Executive Summary

Since its inception the WHO Model Essential Medicines List (EML) has evolved and matured into a critically important element of national health system policies and practices and has proven to be a useful tool to help countries in selecting essential medicines. WHO recommends the selection of essential drugs as one of the core principles of a national drug policy because it helps set priorities for all aspects of the pharmaceutical systems shown in Figure 1.

WHO defines essential medicines as comprising those “that satisfy the needs of the majority of the population” (http://www.who.int/medicines/services/essmedicines_def/en/).



Key Policy Issues:


Drug selection, preferably linked to national clinical guidelines, is a crucial step in ensuring access to essential drugs and in promoting rational drug use, because no public sector or health insurance system can afford to supply or reimburse all drugs that are available on the market. Key policy issues are:

1. the adoption of the essential drugs concept to identify priorities for government involvement in the pharmaceutical sector, and especially for drug supply in the public sector and for reimbursement schemes;
2. procedures to define and update the national list(s) of essential drugs; selection mechanisms for traditional and herbal medicines. Selection mechanisms for traditional and herbal medicines.

Figure 1: Drug Selection

Every country defines its own list of essential medicines in accordance with the health status and requirements of its population. The list is revised periodically to take into account the changing prevalence of new, emerging or re-emerging diseases and new therapeutic developments; ensures drug quality and provides for the continued development of better medicines that also meet changing resistance patterns.

According to WHO Access to Medicine Framework (Figure 2) by virtue of the fact that the majority of the population in developing countries lacks access to essential medicines, therefore the EML can be used as an advocacy tool for governments to draw up their national lists to make medicines accessible, spending their resources on those medicines that are most needed and offer the best value for money.



Significance

Essential medicines are intended to be available within the context of functioning health systems at all times in adequate amounts, in the appropriate dosage forms, with assured quality and at a price the individuals and the community can afford. Essential Medicine List is a tool for national health system policy towards selection of essential medicines and to ensure drug quality; and the need for continued development of better medicines for emerging diseases as well as to meet changing resistance patterns. Proper use of essential medicines improves patient safety and limits unnecessary medical spending. Lists of essential medicines also guide the procurement and supply of medicines in the public sector. WHO recommends revision of EML after every two years. NEML 2018 is being reviewed after the latest NEML that was developed in 2016.

Rationale

NEML 2018 is being developed considering the **WHO Model List 2017**, containing essential medicines with respect to global requirements, and has been customized per local requirements and priorities, Provincial essential medicine priorities and recommendations of healthcare experts.

The criteria for selection of essential medicines was based on disease prevalence, public health relevance, evidence of clinical efficacy and safety, comparative cost-effectiveness and new interventions emerging as a result of research and development in pharmaceutical industry to address the challenges of new diseases emerging globally. This is especially important for Pakistan where availability of medicines in public sector facilities is reported to be 39% (World Health Organization country office; Pakistan, 2014).

The National Essential Medicines (Drugs) List (NEML) of Pakistan was first prepared in 1994 after consultation with experts in the field. The list was reviewed in 1995, 2000, 2003, 2007 & 2013 by Ministry of Health (defunct). After devolution and

promulgation of DRAP (Drug Regulatory Authority of Pakistan) ACT 2012; Pharmacy Services Division was established under the Drug Regulatory Authority of Pakistan and assigned the responsibility to develop and periodically review NEML based on the Pakistan's disease burden, priority health conditions and affordability concerns. In this regard, the last edition of the NEML was published in 2016 which now has been reviewed and current edition of NEML 2018 has been published comprising of 428 molecules under 30 categories and 78 sub-categories.

Explanatory Note

The following symbols are used throughout the National Essential Medicine List of Pakistan 2018.

The NEML 2018 presents a list of minimum medicine needed for a basic healthcare system, listing the most efficacious, safe and cost-effective medicines for priority conditions. Priority conditions are selected on the basis of current and estimated future public health relevance and potential for safe and cost-effective treatment.

The square box symbol (□) is primarily intended to indicate similar clinical performance within a pharmacological class. The listed medicine should be the example of the class for which there is the best evidence for effectiveness and safety. In some cases, this may be the first medicine that is licensed for marketing; in other instances, subsequently licensed compounds may be safer or more effective. Where there is no difference in terms of efficacy and safety data, the listed medicine should be the one that is generally available at the lowest price, based on international drug price information sources. Therapeutic equivalence is indicated only on the basis of reviews of efficacy and safety and when consistent with WHO clinical guidelines.

The [a] symbol indicates that there is an age or weight restriction on use of the medicine; details for each medicine can be found in Table I

The [D] symbol represents that molecules have been listed in duplicate against various therapeutic categories.

The [c] symbol is placed next to an individual medicine or strength of medicine it signifies that there is a specific indication for restricting its use to children.

The (P) symbol is placed next to an individual medicine which was part of the complementary list in the previous edition and now has been included in the core list of NEML, 2018.

The presence of an entry on the Essential Medicines List carries no assurance as to pharmaceutical quality. It is the responsibility of the relevant national or regional drug

regulatory authority to ensure that each product is of appropriate pharmaceutical quality (including stability) and that, when relevant, different products are interchangeable. Medicines and dosage forms are listed in alphabetical order within each section and there is no implication of preference for one form over another. Standard treatment guidelines should be consulted for information on appropriate dosage forms.

The current publication of National Essential Medicine List 2018 is a guide for healthcare professionals to promote and increase the acceptability of concept and envision to promote national health strategies and reduce the disease burden in the country.



Sheikh Ansar Ahmad,
Director,
Pharmacy Services Division,
Drug Regulatory Authority of Pakistan,
Islamabad.

Alphabetical List NEML 2018

- | | | | |
|-----|--|-----|--|
| 1. | abacavir (ABC) | 30. | artemether + lumefantrine |
| 2. | acetazolamide | 31. | artesunate |
| 3. | acetic acid | 32. | artesunate + sulphadoxine
and pyrimethamine |
| 4. | acetylcysteine | 33. | ascorbic acid |
| 5. | acetylsalicylic acid | 34. | asparaginase |
| 6. | aciclovir | 35. | atazanavir |
| 7. | adenosine | 36. | atazanavir + ritonavir |
| 8. | albendazole | 37. | atenolol |
| 9. | alcohol-based hand rub | 38. | atorvastatin |
| 10. | allopurinol | 39. | atracurium |
| 11. | all-trans retinoid acid
(ATRA) | 40. | atropine |
| 12. | aluminium hydroxide +
magnesium trisilicate | 41. | azathioprine |
| 13. | amidotrizoate | 42. | azithromycin |
| 14. | amikacin | 43. | aztreonam |
| 15. | amiloride | 44. | B complex (B1/ B6 and B12) |
| 16. | aminophylline | 45. | barium sulfate |
| 17. | amiodarone | 46. | BCG vaccine |
| 18. | amitriptyline | 47. | beclomethasone |
| 19. | amlodipine | 48. | bedaquiline |
| 20. | amoxicillin | 49. | bendamustine |
| 21. | amoxicillin + clavulanic acid | 50. | benzathine benzylpenicillin |
| 22. | amphotericin B | 51. | benzoyl peroxide |
| 23. | ampicillin | 52. | benzyl benzoate |
| 24. | anastrozole | 53. | benzylpenicillin |
| 25. | anti-D immunoglobulin | 54. | betamethasone |
| 26. | anti-hepatitis B
immunoglobulin | 55. | bevacizumab |
| 27. | anti-rabies immunoglobulin | 56. | bicalutamide |
| 28. | anti-tetanus immunoglobulin | 57. | biperiden |
| 29. | anti-venom immunoglobulin | 58. | bisacodyl |
| | | 59. | bisoprolol |
| | | 60. | bleomycin |

61.	budesonide	100.	cloxacillin + ampicillin
62.	bupivacaine	101.	clozapine
63.	caffeine citrate	102.	coagulation factor VIII
64.	calamine	103.	coagulation factor IX
65.	calcium	104.	coal tar
66.	calcium folinate	105.	codeine
67.	calcium gluconate	106.	condoms
68.	capecitabine	107.	copper-containing device
69.	capreomycin	108.	cyclizine
70.	carbamazepine	109.	cyclophosphamide
71.	carboplatin	110.	cycloserine
72.	cefalexin	111.	cytarabine
73.	cefazolin	112.	dacarbazine
74.	cefixime	113.	daclatasvir
75.	ceftazidime	114.	dactinomycin
76.	ceftriaxone	115.	darunavir
77.	charcoal/ activated	116.	dasabuvir
78.	chlorambucil	117.	daunorubicin
79.	chloramphenicol	118.	deferoxamine
80.	chlorhexidine	119.	delamanid
81.	chlorine base compound	120.	desmopressin
82.	chloroquine	121.	dexamethasone
83.	chloroxylenol	122.	dextran 40
84.	chlorpheniramine	123.	dextran 70
85.	chlorpromazine	124.	diaphragm
86.	cholecalciferol	125.	diazepam
87.	cholera vaccine	126.	diclofenac sodium
88.	ciclosporin	127.	diethylcarbamazine
89.	cinnarizine	128.	digoxin
90.	ciprofloxacin	129.	dihydroartemisinin +
91.	cisplatin		piperquine
92.	clarithromycin	130.	diloxanide
93.	clindamycin	131.	dimenhydrinate
94.	clofazimine	132.	diphtheria antitoxin
95.	clomifene	133.	diphtheria vaccine
96.	clomipramine	134.	dobutamine
97.	clopidogrel	135.	docetaxel
98.	clotrimazole	136.	dolutegravir
99.	cloxacillin	137.	dopamine

138.	doxorubicin	170.	fluorouracil
139.	doxycycline	171.	fluoxetine
140.	efavirenz (EFV or EFZ)	172.	fluphenazine
141.	enalapril	173.	folic acid
142.	enoxaparin	174.	fresh-frozen plasma
143.	entecavir	175.	furosemide
144.	ephedrine	176.	gemcitabine
145.	ephedrine	177.	gentamicin
146.	epinephrine (adrenaline)	178.	glibenclamide
147.	ergocaliferol	179.	gliclazide
148.	ergometrine	180.	glucagon
149.	erythromycin	181.	glucose
150.	estradiol cypionate + medroxyprogesterone acetate	182.	glucose with sodium chloride
151.	estrogen	183.	glutaral
152.	ethambutol	184.	glycerin
153.	ethambutol + isoniazid	185.	glyceryl trinitrate
154.	ethambutol + isoniazid + pyrazinamide + rifampicin	186.	glycopyrrolate
155.	ethambutol + isoniazid + rifampicin	187.	griseofulvin
156.	ethanol	188.	Haemophilus influenzae type b vaccine
157.	ethinylestradiol + levonorgestrel	189.	haloperidol
158.	ethinylestradiol + norethisterone	190.	heparin sodium
159.	ethionamide	191.	hepatitis A vaccine
160.	etongestrel-releasing implant	192.	hepatitis B vaccine
161.	etoposide	193.	HPV vaccine
162.	ferrous salt	194.	hydralazine
163.	ferrous salt + folic acid	195.	hydrochlorothiazide
164.	filgrastim	196.	hydrocortisone
165.	fluconazole	197.	hydroxocobalamin
166.	flucytosine	198.	hydroxycarbamide
167.	fludarabine	199.	hydroxychloroquine
168.	fludrocortisone	200.	ibuprofen
169.	fluorescein	201.	ifosfamide
		202.	imatinib
		203.	influenza vaccine (seasonal)
		204.	insulin injection (soluble)
		205.	intermediate-acting insulin
		206.	Intraperitoneal dialysis solution

207.	iodine	241.	lugol's solution
208.	iohexol	242.	maggot oil (turpentine oil)
209.	ipratropium bromide	243.	magnesium oxides and hydroxides
210.	irinotecan	244.	magnesium sulfate
211.	iron sucrose	245.	mannitol
212.	isoflurane	246.	measles vaccine
213.	isoniazid	247.	mebendazole
214.	isoniazid + pyrazinamide + rifampicin	248.	medroxyprogesterone acetate
215.	isoniazid + rifampicin	249.	mefloquine
216.	isoniazid+pyridoxine+sulfame thoxazole+trimethoprim	250.	meglumine antimoniate
217.	isoprenaline	251.	meglumine iotroxate
218.	isosorbide dinitrate	252.	meningococcal meningitis vaccine
219.	itraconazole	253.	mercaptopurine
220.	ivermectin	254.	meropenem
221.	kanamycin	255.	mesna
222.	ketamine	256.	metformin
223.	labetalol	257.	methadone
224.	lamivudine (3TC)	258.	methotrexate
225.	latanoprost	259.	methylated spirit
226.	ledipasvir + sofosbuvir	260.	methyldopa
227.	leuprorelin	261.	methylprednisolone
228.	levamisole	262.	methylthioninium chloride (methylene blue)
229.	levofloxacin	263.	metoclopramide
230.	levonorgestrel	264.	metronidazole
231.	levonorgestrel-releasing implant	265.	miconazole
232.	levonorgestrel-releasing intrauterine system	266.	midazolam
233.	levothyroxine	267.	miltefosine
234.	lidocaine	268.	misoprostol
235.	lidocaine + epinephrine (adrenaline)	269.	montelukast
236.	linezolid	270.	morphine
237.	lithium carbonate	271.	moxifloxacin
238.	loperamide	272.	multiple micronutrients
239.	loratadine	273.	multivitamins
240.	losartan potassium	274.	mumps vaccine
		275.	mupirocin

276.	nalbuphine HCL	312.	phenytoin
277.	naloxone	313.	phloroglucinol
278.	naproxen	314.	phytomenadione
279.	natamycin	315.	pilocarpine
280.	neostigmine	316.	piperacillin + tazobactum
281.	nevirapine (NVP)	317.	platelets
282.	nicotinamide	318.	pneumococcal vaccine
283.	nicotine replacement therapy (NRT)	319.	podophyllum resin
284.	nifedipine	320.	poliomyelitis vaccine
285.	nitrofurantoin	321.	polymyxin B (sulphate)+ bacitracin zinc
286.	nitrous oxide	322.	potassium chloride
287.	norethisterone enantate	323.	potassium iodide
288.	normal immunoglobulin	324.	potassium permanganate
289.	nystatin	325.	povidone iodine
290.	ofloxacin	326.	praziquantel
291.	olanzapine	327.	prednisolone
292.	ombitasvir + paritaprevir + ritonavir	328.	pregabalin
293.	omeprazole	329.	primaquine
294.	ondansetron	330.	procaine benzyl penicillin
295.	oral rehydration salts	331.	procarbazine
296.	oxaliplatin	332.	progesterone vaginal ring
297.	oxygen	333.	proguanil
298.	oxytocin	334.	proparacaine (hydrochloride)
299.	paclitaxel	335.	propofol
300.	p-aminosalicylic acid	336.	propranolol
301.	pancreatic enzymes	337.	propylthiouracil
302.	pancuronium	338.	prostaglandin E
303.	paracetamol	339.	protamine sulfate
304.	paromomycin	340.	pyrantel
305.	pegylated interferon alfa (2a or 2b)	341.	pyrazinamide
306.	penicillamine	342.	pyridostigmine
307.	permethrin	343.	pyridoxine
308.	pertussis vaccine	344.	rabies vaccine
309.	pheniramine	345.	raltegravir
310.	phenobarbital	346.	ranitidine
311.	phenoxymethylpenicillin	347.	red blood cells
		348.	retinol
		349.	ribavirin

350.	riboflavin	386.	sulfadoxine + pyrimethamine
351.	rifabutin		co-blister / combined
352.	rifampicin		therapy (For Falciparum)
353.	rifapentine	387.	sulfamethoxazole +
354.	risovastatin		trimethoprim
355.	risperidone	388.	sulfasalazine
356.	ritonavir	389.	sumatriptan
357.	rituximab	390.	surfactant
358.	rotavirus vaccine	391.	suxamethonium
359.	rubella vaccine	392.	tamoxifen
360.	salbutamol	393.	tears natural
361.	salicylic acid	394.	tenofovir disoproxil fumarate
362.	salicylic acid + betamethasone	395.	(TDF) terbinafine
363.	selenium sulfide	396.	testosterone
364.	senna	397.	tetanus immunoglobulin
365.	sevoflurane		(human)
366.	silver sulfadiazine	398.	tetanus vaccine
367.	simeprevir	399.	tetracaine
368.	simvastatin	400.	tetracycline
369.	sodium biphosphate	401.	thiamine
370.	sodium chloride	402.	thioguanine
371.	sodium cromoglycate	403.	timolol
372.	sodium fluoride	404.	tirofiban
373.	sodium hydrogen carbonate	405.	tobramycin + dexamethasone
374.	sodium lactate/ compound solution	406.	tramadol
375.	sodium nitrite	407.	tranexamic acid
376.	sodium nitroprusside	408.	trastuzumab
377.	sodium polystyrene sulfonate	409.	tropicamide
378.	sodium stibogluconate	410.	tuberculin/ purified protein derivative (PPD)
379.	sodium thiosulfate	411.	typhoid vaccine
380.	sodium thiosulphate	412.	urea
381.	sofosbuvir	413.	valproic acid (sodium valproate) (D)
382.	spectinomycin	414.	vancomycin
383.	spironolactone	415.	varicella vaccine
384.	streptokinase	416.	vecuronium
385.	streptomycin		

- 417. verapamil
- 418. vinblastine
- 419. vincristine
- 420. vinorelbine
- 421. vitamin K1 (phytonadione)
- 422. warfarin (D)
- 423. water for injection
- 424. whole blood
- 425. xylocaine
- 426. yellow fever vaccine
- 427. zidovudine (ZDV or AZT).
- 428. zinc sulfate

National Essential Medicine List of Pakistan 2018

Therapeutic Category	Generic Name	Formulations	Service Level Category
I. Anesthetics			
I.1 General anesthetics and oxygen			
I.1.1 Inhalational medicines	isoflurane sevoflurane (P) nitrous oxide oxygen ketamine propofol glycopyrrolate (P)	inhalation inhalation inhalation inhalation (medicinal gas) injection containing ketamine hydrochloride: 50 mg/ml in 10-ml vial injection: 10 mg/ml; 20 mg/ml Injection: 0.2 mg/ml	S, T T P, S, T P, S, T P, S, T P, S, T S, T
I.1.2 Injectable medicines			
I.2 Local anesthetics			
	<input type="checkbox"/> bupivacaine	injection: 0.25%; 0.5% (hydrochloride) in vial injection for spinal anesthesia: 0.5% (hydrochloride) in 4-ml ampoule to be mixed with 7.5% glucose solution	S, T S, T
	<input type="checkbox"/> lidocaine (D)	injection: 1%; 2% (hydrochloride) in vial injection for spinal anesthesia: 5% (hydrochloride) in 2-ml ampoule mixed with 7.5% glucose solution	P, S, T P, S, T
		gel lidocaine HCl 2% w/v 15 gm tube	P, S, T
		dental cartridge: 2% (hydrochloride) + epinephrine 1:80 000 injection: 1%; 2% (hydrochloride or sulfate) + epinephrine 1:200 000 in vial	P, S, T
	ephedrine	injection: 30 mg (hydrochloride)/ml in 1-ml ampoule (for use in spinal anesthesia during delivery, to prevent hypotension)	P, S, T S, T
I.3 Preoperative medication and sedation for short-term procedures			
	atropine (D)	injection: 1 mg (sulfate) in 1-ml ampoule	P, S, T
	<input type="checkbox"/> midazolam	injection: 1 mg/ml oral liquid: 2 mg/ml [c] tablet: 7.5 mg; 15 mg	P, S, T P, S, T P, S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
2. Medicines for Pain and Palliative Care			
2.1. Non-opioids analgesics, antipyretics, and non-steroidal anti-inflammatory medicine (NSAIM)			
	acetylsalicylic acid (D)	suppository: 50 mg to 150 mg tablet: aspirin 75 mg enteric coated tablet/blister suspension: ibuprofen 100 mg/5 ml bottle tablet: ibuprofen 400 mg blister <i>(not for children less than 3 months old)</i>	P, S, T
	ibuprofen (a) (D)	syrup: paracetamol 120 mg/5 ml bottle suppository: 100 mg	P, S, T
	paracetamol(D)	injection: 150 mg/ml tablet: paracetamol 500 mg blister <i>*not recommended for anti-inflammatory use; has no proven benefit</i>	P, S, T
	diclofenac sodium	tablet: 50 mg	P, S, T
	naproxen	injection: 75 mg/5 ml tablets:250, 500 mg	S, T
2.2. Opioid analgesics			
	codeine (D)	tablet: 30 mg (phosphate)	T
	nalbuphine HCl	injection: 10mg/ml granules (slow release; mix with water): 20 mg–200 mg (morphine sulfate)	T
	<input type="checkbox"/> morphine (D)*	injection: 10 mg (morphine hydrochloride or morphine sulphate in 1 ml ampoule) oral liquid: 10 mg (morphine hydrochloride or morphine sulfate)/5 ml tablet (slow release): 10 mg–200 mg (morphine hydrochloride or morphine sulfate) tablet (immediate release): 10 mg (morphine sulfate) <i>*Alternatives limited to hydromorphone and oxycodone.</i>	S, T
2.3. Medicines for other common symptoms in palliative care			
	amitriptyline	tablet: amitriptyline HCl 10 mg, 25 mg	T
	cyclizine	injection: 50 mg/ml tablet: 50 mg	P, S, T
	dexamethasone (D)	injection: 4 mg/ml in 1-ml ampoule (as disodium phosphate salt) oral liquid: 2 mg/5 ml	P, S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
		tablet: 2 mg [C]; 4 mg injection: 5 mg/ml oral liquid: 2 mg/5 ml rectal solution: 2.5 mg, 5 mg, 10 mg tablet: 5 mg, 10 mg injection: 5 mg in 1-ml ampoule oral liquid: 2 mg/ml solid oral dosage form: 0.5 mg, 2 mg, 5 mg solid oral dosage form: 2 mg injection: 5 mg (hydrochloride)/ml in 2-ml ampoule oral liquid: 5 mg/5 ml solid oral form: 10 mg (hydrochloride) injection: 2 mg base/ml in 2-ml ampoule (as hydrochloride)	P, S, T P, S, T P, S, T S, T S, T S, T P, S, T S, T S, T S, T S, T S, T
	diazepam (D)	oral liquid: 2.5 mg, 5 mg, 10 mg tablet: 5 mg, 10 mg injection: 5 mg in 1-ml ampoule oral liquid: 2 mg/ml solid oral dosage form: 0.5 mg, 2 mg, 5 mg solid oral dosage form: 2 mg injection: 5 mg (hydrochloride)/ml in 2-ml ampoule oral liquid: 5 mg/5 ml solid oral form: 10 mg (hydrochloride) injection: 2 mg base/ml in 2-ml ampoule (as hydrochloride)	P, S, T P, S, T S, T S, T S, T S, T P, S, T S, T S, T S, T S, T S, T
	haloperidol (D)		
	loperamide		
	metoclopramide		
	ondansetron [C] (a)	solid oral dosage form: equivalent to 4 mg base; equivalent to 8 mg base; (a) >/ month tablet: 5 mg capsules: 75 mg, 150 mg, 300 mg capsules: 50 mg. injection: 50 mg/ml.	S, T P, S, T S, T S, T S, T S, T S, T
	bisacodyl		
	pregabalin (P)		
	tramadol (P)		
3. Anti-Allergics and Medicines Used in Anaphylaxis			
	chlorpheniramine	tablets: 4 mg (hydrogen maleate)	P, S, T
	dexamethasone (D)	injection: 10 mg/ml (hydrogen maleate in 1-ml ampoule)	P, S, T
	epinephrine (adrenaline) (D)	injection: 4 mg/ml in 1-ml ampoule (as disodium phosphate salt)	P, S, T
	hydrocortisone (D)	injection: 1 mg (as hydrochloride or hydrogen tartrate) in 1-ml ampoule powder for injection: 100 mg, 250 mg, 1 gram (as sodium succinate) in vial oral liquid: 1 mg/ml	P, S, T P, S, T
	□ loratadine *	tablet: 10 mg	P, S, T
	□ prednisolone (D)	*There may be a role for sedating antihistamines for limited indications (EMIC)	P, S, T
		oral liquid: 5 mg/ml [C] tablet: 5 mg, 25 mg	P, S, T

Therapeutic Category	Generic Name		Formulations	Service Level Category
	pheniramine (P)		injection: pheniramine (maleate): 22.7 mg/ml in 2 ml ampoule equivalent to 25 mg pheniramine salicylate. tablet: 25 mg, 50 mg. elixir/syrup: 15 mg/5 ml.	P, S, T
				P, S, T
				P, S, T
4. Antidotes and Other Substances Used in Poisoning				
4.1. Non-specific	charcoal, activated	powder		P, S, T
4.2. Specific				
	acetylcysteine	injection: 200 mg/ml in 10-ml ampoule oral liquid: 10% [c]; 20% [c]		P, S, T
	atropine	injection: 1 mg (sulfate) in 1-ml ampoule		P, S, T
	calcium gluconate (D)	injection: 100 mg/ml in 10-ml ampoule		P, S, T
	methylthioninium chloride (methylene blue)	injection: 10 mg/ml in 10-ml ampoule		S, T
	naloxone	injection: 400 micrograms (hydrochloride) in 1-ml ampoule		P, S, T
	sodium nitrite	injection: 30 mg/ml in 10-ml ampoule		S, T
	deferoxamine	powder for injection: 500 mg (mesilate) in vial		S, T
5. Anticonvulsants/Antiepileptics				
	carbamazepine (D)	oral liquid: 100 mg/5 ml tablet (chewable): 100 mg, 200 mg tablet (scored): 100 mg, 200 mg		P, S, T
	diazepam (D)	gel or rectal solution: 5 mg/ml in 0.5-ml, 2-ml, 4-ml tubes		P, S, T
	magnesium sulfate*	injection: 0.5g/ml in 2-ml ampoule (equivalent to 1 gram in 2-ml; 50% weight/volume); 0.5g/ml in 10-ml ampoule (equivalent to 5 grams in 10-ml; 50% weight/volume)		P, S, T
		*For use in eclampsia and severe pre-eclampsia, but not for other convulsant disorders.		
	phenobarbital	injection: 200 mg/ml (sodium) oral liquid: 15 mg/5 ml tablet: 15 mg to 100 mg		P, S, T
				P, S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
		injection: 50 mg/ml in 5-ml vial (sodium salt) oral liquid: 25 mg to 30 mg/5 ml* solid oral dosage form: 25 mg, 50 mg, 100 mg (sodium salt)	P, S, T
	phenytoin	tablet (chewable): 50 mg *having both 25 mg/5 ml and 30 mg/5 ml strengths in the same market would be confusing in prescribing and dispensing and should be avoided	P, S, T
		oral liquid: 200 mg/5 ml tablet (enteric coated): 200 mg, 500 mg as sodium valproate	P, S, T
	valproic acid (sodium valproate) (D)	injection: 100 mg/ml in 4-ml ampoule; 100 mg/ml in 10-ml ampoule	T
6. Anti-Infective Medicines			
6.1. Anthelmintics			
6.1.1 Intestinal anthelmintics	albendazole (D) mebendazole pyrantel ivermectin levamisole diethylcarbamazine	tablet (chewable): 400 mg oral liquid: 100 mg/5ml tablet (chewable): 100 mg, 500 mg oral liquid: 50 mg (as embonate or pamoate)/ml tablet (chewable): 250 mg (as embonate or pamoate) tablet: 3 mg tablet: 50 mg; 150 mg (as hydrochloride) tablet: 50 mg, 100 mg (dihydrogen citrate)	P, S, T
6.1.2 Antifilarials			P, S, T
6.1.3 Antischistosomals and other antirematode medicines	praziquantel (D)	tablet: 150 mg, 600 mg	S, T
6.2. Antibacterials			
6.2.1 Beta-lactam medicines			
	amoxicillin	powder for oral liquid: 125 mg (as trihydrate)/5 ml; 250 mg (as trihydrate)/5 ml [c] injection 250 mg, 500 mg	P, S, T
			P, S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
	amoxicillin + clavulanic acid	solid oral dosage form: 250 mg, 500 mg (as trihydrate) oral liquid: 125 mg amoxicillin + 31.25 mg clavulanic acid/5 ml and 250 mg amoxicillin + 62.5 mg clavulanic acid/5 ml [c] tablet: 500 mg (as trihydrate) + 125 mg (as potassium salt)	P, S, T S, T
	ampicillin	powder for injection: 500 mg; 1 gram (as sodium salt) in vial capsules: 250 mg, 500 mg syrups: 125 mg, 250 mg/ml	P, S, T P, S, T P, S, T
	benzathine benzylpenicillin	powder for injection: 900 mg benzylpenicillin (= 1.2 million IU) in 5-ml vial [c]; 1.44 g benzylpenicillin (= 2.4 million IU) in 5-ml vial	P, S, T
	benzylpenicillin	powder for injection: 600 mg (= 1 million IU); 3 grams (= 5 million IU) (sodium or potassium salt) in vial	P, S, T
	phenoxyymethylpenicillin	powder for oral liquid: 250 mg (as potassium salt)/5 ml tablet: 250 mg (as potassium salt)	P, S, T P, S, T
	procaine benzyl penicillin*	powder for injection: 1 gram (= 1 million IU); 3 grams (= 3 million IU) in vial *procaine benzylpenicillin is not recommended as first-line treatment for neonatal sepsis, except in settings with high neonatal mortality, and when given by trained health workers if hospital care is not possible	P, S, T P, S, T
	cefalexin [c]	powder for reconstitution with water: 125 mg/5 ml, 250 mg/5 ml (anhydrous)	P, S, T
	□ cefazolin (a)*	solid oral dosage form: 250 mg (as monohydrate) powder for injection: 1 g (as sodium salt) in vial *for surgical prophylaxis. (a) > 1 month.	P, S, T P, S, T
	□ cloxacillin	capsule: 500 mg; 1 gram (as sodium salt) powder for injection: 500 mg (as sodium salt) in vial	P, S, T P, S, T
	cloxacillin + ampicillin	powder for suspension ©: 125 mg & 250 mg /5ml	S, T
Watch Group Antibiotics			
	ceftriaxone (a)*	powder for injection: 250 mg, 500 mg; 1 gram (as sodium salt) in vial *do not administer with calcium and avoid in infants with hyperbilirubinaemia. (a) >4 weeks corrected gestational age	S, T
	cefixime*	capsule: 400 mg (as trihydrate) *listed only for single-dose treatment of uncomplicated <u>anogenital gonorrhoea</u>	S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
		suspension: 100 mg/5 ml; 200 mg/5 ml. (a) >4 / weeks corrected gestational age Powder for injection 2 g (as sodium salt) + 250 mg (as sodium salt) ; 4 g (as sodium salt) + 500 mg (as sodium salt) in vial powder for injection: 250 mg/500 mg/1 gram (as pentahydrate) in vial injection: 500 mg in vial	S, T
	piperacillin + tazobactum		
	ceftazidime		
	meropenem		
6.2.2 Other antibacterials			
Key Access Antibiotics			
	chloramphenicol	capsule 250 mg oral liquid 150 mg (as palmitate)/ 5 ml powder for injection 1 g (sodium succinate) in vial	T T T
	amikacin (sulphate) (Reserved 2nd line)	injection: 100mg; 250mg capsule: 150 mg/300 mg (as hydrochloride) injection: 150 mg (as Phosphate)/ml	T S, T
	clindamycin	oral liquid: 75 mg/5 ml (as palmitate) [C] oral liquid: 25 mg/5 ml [C]; 50 mg/5 ml (anhydrous) [C]	S, T P, S, T
	doxycycline (a) (D)	solid oral dosage form: 50 mg [C]; 100 mg (as hydclate) (a) use in children <8 years only for life-threatening infections when there is no alternative	P, S, T P, S, T
		oral liquid: 250 mg/5 ml (anhydrous) [C]	S, T
		solution for iv infusion: 2 mg/ml (as hydclate) [C]	S, T
		tablet: coated, 250 mg/500 mg (as hydrochloride)	S, T
		*square box is for adults only	
	moxifloxacin (P)	tablet: 200 mg/400 mg.	S, T
Watch Group Antibiotics			
	azithromycin*	capsule: 250 mg; 500 mg (anhydrous) oral liquid: 200 mg/5 ml	P, S, T
		*only listed for single-dose treatment of <i>genital chlamydia trachomatis</i> and trachoma	
	clarithromycin*	oral liquid: 125 mg/5ml solid oral dosage form: 500 mg	S, T
		*for use in combination regimens to eradicate <i>H. pylori</i> in adults	S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
	doxycycline (a) (D)	oral liquid: 25 mg/5 ml [c]; 50 mg/5 ml (anhydrous) [c] solid oral dosage form: 50 mg [c]; 100 mg (as hydclate) (a) use in children <8 years only for life-threatening infections when there is no alternative	P, S, T
	<input type="checkbox"/> gentamicin (D)	injection: 10 mg, 40 mg (as sulfate)/ml in 2-ml vial	P, S, T
	<input type="checkbox"/> metronidazole (D)	injection: 500 mg in 100-ml vial oral liquid: 200 mg (as benzoate)/5 ml	P, S, T
	<input type="checkbox"/> nitrofurantoin	tablet: 400 mg oral liquid: 25 mg/5 ml [c] tablet: 100 mg	P, S, T
		injection: 80 mg + 16 mg/ml in 5-ml ampoule, 80 mg + 16 mg/ml in 10-ml ampoule oral liquid: 200 mg + 40 mg/5ml	P, S, T
		tablet: 100 mg + 20 mg, 400 mg + 80 mg, 800 mg + 160 mg	P, S, T
	specinomycin	powder for injection 2 g (as hydrochloride) in vial	T
	vancomycin	powder for injection: 250 mg/500 mg (as hydrochloride) in vial	T
	aztreonam (P)	injection: contains aztreonam 500 mg, 1g vial.	T
6.2.3 Antituberculosis medicines	WHO recommends and endorses the use of fixed-dose combinations and the development of appropriate new fixed-dose combinations, including modified dosage forms, non-refrigerated products, and pediatric dosage forms of assured pharmaceutical quality.		
	ethambutol	oral liquid: 25 mg/ml [c] tablet: 100 mg to 400 mg (hydrochloride)	
	ethambutol + isoniazid	tablet: 400 mg + 150 mg	
	ethambutol + isoniazid + pyrazinamide + rifampicin	tablet: 275 mg + 75 mg + 400 mg + 150 mg	
	ethambutol + isoniazid + rifampicin	tablet: 275 mg + 75 mg + 150 mg	
	isoniazid	oral liquid: 50 mg/5 ml [c] tablet: 100 mg to 300 mg tablet (scored): 50, mg	
	isoniazid + pyrazinamide + rifampicin	tablet: 75 mg + 400 mg + 150 mg, 150 mg + 500 mg + 150 mg (for intermittent use three times weekly)	
	isoniazid + rifampicin	tablet: 75 mg + 150 mg, 150 mg + 300 mg 60 mg + 60 mg (for intermittent use three times a week)	

Therapeutic Category	Generic Name	Formulations	Service Level Category
		150 mg + 150 mg (for intermittent use three times a week) oral liquid: 30 mg/ml [c]	
	pyrazinamide	tablet: 400 mg tablet (dispersible): 150 mg tablet (scored): 150 mg capsule: 150 mg.*	
	rifabutin	*only use in patients who are receiving protease inhibitors.	
	rifampicin (D)	oral liquid: 20 mg/ml [c] solid oral dosage form: 150 mg, 300 mg tablet: 150 mg	
	rifapentine*	*for treatment of latent TB infection (LTBI) only	
	streptomycin	powder for injection: 1 gram (as sulfate) in vial	
	amikacin	powder for injection: 100 mg, 500 mg, 1 gram (as sulfate) in vial	S, T
	bedaquiline	tablet: 100 mg	
	capreomycin	powder for injection: 1 gram (as sulfate) in vial	
	cycloserine	solid oral dosage form: 250 mg	
	delamanid	*can use terizidone as an alternative tablet: 50 mg	
	ethionamide*	tablet: 125 mg, 250 mg *can use prafloxacinamide as an alternative	
	kanamycin	powder for injection: 1 gram (as sulfate) in vial	
	levofloxacin*	tablet: 250 mg/500 mg/750 mg *can use ofloxacin and moxifloxacin as alternatives, depending on availability and program considerations	S, T
	linezolid	injection for intravenous administration: 2 mg/ml in 300 ml bag powder for oral liquid: 100 mg/5 ml	
	P-aminosalicylic acid	tablet: 400 mg, 600 mg granules: 4 grams in sachet	As per NTP guidelines.
	clofazimine	tablet: 500 mg capsules: 50mg; 100mg	

Therapeutic Category	Generic Name	Formulations	Service Level Category
6.3. Antifungal medicines			
	amphotericin B (D)	powder for injection: 50 mg in vial (as sodium deoxycholate or liposomal complex)	S, T
	clotrimazole	vaginal cream: 1%, 10% vaginal tablet: 100 mg, 500 mg with applicator	P, S, T
	<input checked="" type="checkbox"/> fluconazole*	capsule: 50 mg, 150 mg, 200 mg injection: 2 mg/ml in vial oral liquid: 50 mg/5 ml capsule: 250 mg infusion: 2.5 grams in 250 ml oral liquid: 125 mg/5 ml [C] solid oral dosage form: 125 mg, 250 mg	P, S, T S, T
	flucytosine	lozenge: 100 000 IU oral liquid: 50 mg/5 ml [C]; 100 000 IU/ml [C]	P, S, T
	griseofulvin	pessary: 100 000 IU tablet: 100 000 IU; 500 000 IU	P, S, T
	nystatin	capsule: 100 mg oral liquid: 10 mg/ ml	P, S, T
	itraconazole		S, T
			S, T
6.4. Antiviral medicines			
		Based on current evidence and experience, medicines in the following three classes of antiretrovirals are included as essential medicines for treatment and prevention of HIV (prevention of mother-to-child transmission and post-exposure prophylaxis). WHO emphasizes the importance of following the global and national guidelines for these products. WHO recommends and endorses the use of fixed dose combinations (FDCs) and the development of appropriate new FDCs, including modified dosage forms, non-refrigerated products, and pediatric dosage forms of assured pharmaceutical quality.	
		Scored tablets can be used for children; they can be included in the list of tablets, if adequate quality products are available.	
6.4.1 Anti-herpes medicines	<input checked="" type="checkbox"/> aciclovir [C] (D)	oral liquid: 200 mg/5 ml powder for injection: 250 vial (as sodium salt) in vial	P, S, T
		tablet: 200 mg	P, S, T
6.4.2.1 Nucleoside/nucleotide reverse	abacavir (ABC) lamivudine (3TC)	oral liquid: 100 mg as sulfate/5 ml oral liquid: 50 mg/5 ml tablet: 150 mg	S, T S, T S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
transcriptase inhibitors	entecavir	tablet:0.5 mg, 1 mg capsule: 100 mg oral liquid: 50 mg/5 ml solution for iv infusion: 10 mg /ml in 20 ml vial	S, T
6.4.2.2 Non-nucleoside reverse transcriptase inhibitors	efavirenz (EFV or EFZ) (a)	tablet: 300 mg tablet: 200 mg (scored), 600 mg (a) >3 years or >10 kg weight oral liquid: 50 mg/5 ml	S, T
	nevirapine (NVP)	tablet: 50 mg (dispersible), 200 mg	S, T
	atazanavir (a)	solid oral dosage form: 100 mg, 150 mg, 300 mg (as sulfate). (a) >25 kg	S, T
	atazanavir + ritonavir	tablet:300 mg (as sulphate) + 100 mg	S, T
6.4.2.3 Protease inhibitors	darunavir (a)	tablet: 75 mg, 400 mg, 600 mg, 800 mg (a) >3 years	S, T
	ritonavir*	oral liquid: 400 mg/5 ml *Ritonavir is recommended for use in combination as a pharmacological booster, not as an antiretroviral. All other protease inhibitors should be used in boosted forms (e.g., with ritonavir).	S, T
	dolutegravir	tablet (heat stable): 25 mg, 100 mg	S, T
	raltegravir	tablet: 50 mg	T
	isoniazid+pyridoxine+sulfamethoxazol e+trimethoprim	tablet: 400 mg	T
6.4.3 Other antivirals	ribavirin*	tablet: 300 mg+ 25 mg+800 mg+160 mg injection for intravenous administration: 800 mg and 1 gram in 10-ml phosphate buffer solution solid oral dosage form: 200 mg, 400 mg, 600 mg	S, T
		*For treatment of viral hemorrhagic fevers.	T
6.4.4 Antihepatitis medicines			
6.4.4.1 Medicines for hepatitis B	entecavir	oral liquid: 0.05 mg/ml	S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
6.4.1.1 Nucleoside/nucleotide reverse transcriptase inhibitors	tenofovir disoproxil fumarate (TDF)	tablet: 300 mg (tenofovir disoproxil fumarate—equivalent to 245 mg tenofovir disoproxil)	S, T
6.4.1.2 Medicines for hepatitis C	<i>Based on current evidence, medicines in the following classes of direct acting antiviral medicines are included as essential medicines for treatment of hepatitis C virus infections. WHO guidelines recommend specific combination therapy using medicines from different classes.</i>		
6.4.2.1 Nucleotide polymerase inhibitors	sofosbuvir	tablet: 400 mg	S, T
6.4.2.2 Protease inhibitors	simeprevir	capsule: 150 mg	S, T
6.4.2.3 NSSA inhibitors	daclatasvir	tablet: 30 mg; 60 mg (as hydrochloride)	S, T
6.4.2.4 Non-nucleoside polymerase inhibitors	dasabuvir	tablet: 250 mg	S, T
	ribavirin*	injection for intravenous administration: 800 mg and 1 gram in 10-ml phosphate buffer solution solid oral dosage form: 200 mg, 400 mg, 600 mg	S, T
6.4.2.5 Other antivirals	pegylated interferon alfa (2a or 2b) * *To treat hepatitis C, in combination with peginterferon and/or direct acting antiviral medicines. vial or prefilled syringe: 180 micrograms (peginterferon alfa-2a), 80 micrograms, 100 micrograms (peginterferon alfa-2b) *Use in combination with ribavirin. ledipasvir + sofosbuvir tablet: 90 mg + 400 mg ombitasvir + paritaprevir + ritonavir tablet: 12.5 mg + 75 mg + 50 mg		

Therapeutic Category	Generic Name	Formulations	Service Level Category	
6.5. Antiprotozoal medicines				
6.5.1 Antiamoebic and antigiardiasis medicines	diloxanide (a) □ metronidazole (D)	tablet: 500 mg (furoate); (a) >25 kg inject: 500 mg in 100-mL vial oral liquid: 200 mg (as benzoate)/5 mL tablet: 400 mg powder for injection: 50 mg in vial (as sodium deoxycholate or liposomal complex)	P, S, T P, S, T P, S, T S, T	
6.5.2 Anti-leishmaniasis medicines	amphotericin B (D) miltefosine paromomycin* meglumine antimoniate sodium stibogluconate	solid oral dosage form: 10 mg/50 mg solution for intramuscular injection: 750 mg of paromomycin base (as the sulfate) injection 30%, equivalent to approximately 8.1% amitriptyline (pentavalent) in 5-mL ampoule injection: 100 mg/mL, 1 vial = 30 mL	P, S, T P, S, T P, S, T S, T S, T	
		Use medicines in combination to treat <i>P. falciparum</i> malaria cases. The list currently recommends combinations based on treatment guidelines. WHO recognizes that not all the FDCs are listed in the WHO treatment guidelines. They encourage the development and rigorous testing for new FDCs. WHO also encourages the development and testing of rectal dosage formulations.		
		tablets: 20 mg + 120 mg [c] tablets (dispersible): 20 mg + 120 mg [c] artemether + lumefantrine* tablets: 20 mg + 120 mg [c]	*Not recommended in the first trimester of pregnancy or in children below 5 kg. Injection: ampoules, containing 60 mg anhydrous artesunic acid with a separate ampoule of 5% sodium bicarbonate solution. Use to manage severe malaria. Rectal dosage form: 50 mg [c], 200 mg capsules (for pre-referral treatment of severe malaria only; take patients to an appropriate health facility for follow-up care) [c]. tablet: 20 mg *Use in combination with either armodiaquine, mefloquine, or sulfadoxine + pyrimethamine. tablet: 7.5 mg, 15 mg (as diphasphosphate).	As per malaria control program guidelines.
		primaquine* tablets: 14 days.	*Use only to achieve radical cure of <i>P. vivax</i> and <i>P. ovale</i> infections; give for 14 days.	

Therapeutic Category	Generic Name	Formulations	Service Level Category
	artesunate + sulphadoxine and pyrimethamine*	co-blister of (6 + 2) tablets, 2 large tablets; each large tablet contains sulphadoxine 500 mg U.S.P. + pyrimethamine 25 mg U.S.P. and 6 small tablets. Each small tablet contains artesunate 50 mg U.S.P.	
	sulfadoxine + pyrimethamine co-blister / combined therapy (For Falciparum) (P)	co-blister of (6 + 3) tablets, 3 large tablets, each contains sulfadoxine 500 mg U.S.P. + pyrimethamine 25 mg U.S.P.; 6 small tablets; each tablet contains artesunate 100 mg U.S.P. tablets 500mg + 25 mg	
	chloroquine (D)*	oral liquid: 50 mg (as phosphate or sulfate)/5 ml tablet: 150 mg (as phosphate or sulfate)	P, S, T
	doxycycline (a)	*For use only in central American regions, for <i>P. Vivax</i> infections.	P, S, T
6.5.3.2 For prophylaxis	mefloquine (a) (D)	solid oral dosage form: 100 mg (as hydrochloride or hydrate) (a) >8 years. tablet: 250 mg (as hydrochloride)	P, S, T
	proguanil*	(a) >5 kg or >3 months tablet: 100 mg (as hydrochloride)	As per malaria control program guidelines.
	dihydroartemisinin + piperaquine (P)	*Use only in combination with chloroquine. film-coated tablet: contains piperazine tetrabiphosphate (as tetrahydrate + dihydroartemisinin); 320 + 40 mg and 160 mg + 20 mg.	
7. Antimigraine Medicines			
7.1. For treatment of acute attack			
	acetilsalicylic acid	tablet: 300 mg to 500 mg	P, S, T
	ibuprofen [c] (D)	tablet: 200 mg, 400 mg	P, S, T
	paracetamol	oral liquid: 125 mg/5 ml [c]	P, S, T
	cinnarizine (P)	tablet: 300 mg to 500 mg tablets: contains cinnarizine 25 mg	P, S, T
	sumatriptan (P)	injection: 6 mg tablets: 50 mg.	S, T
7.2. For prophylaxis			
	<input type="checkbox"/> propranolol	tablet: 20 mg, 40 mg (hydrochloride)	P, S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
8. Anti-Neoplastic and Immunosuppressives			
Use the medicines listed below only according to the protocols for treating the diseases.			
8.1. Immunosuppressive medicines			
	azathioprine (D)	powder for injection: 100 mg (as sodium salt) in vial tablet (scored): 50 mg capsule: 25 mg	T
	ciclosporin	concentrate for injection: 50 mg/ml in 1-ml ampoule for organ transplantation	T
8.2. Cytoxic and adjuvant medicines			
	all-trans retinoid acid (ATRA)	capsule: 10 mg o- acute promyelocytic leukemia.	T
	allopurinol [C] (D)	tablet: 100 mg, 300 mg powder for injection: 10 000 IU in vial	S, T
	asparaginase	o- acute lymphoblastic leukemia. injection: 45 mg/0.5 ml, 180 mg/2 ml o- chronic lymphocytic leukemia -follicular lymphoma.	T
	bendamustine	powder for injection: 15 mg (as sulfate) in vial -Hodgkin lymphoma	T
	bleomycin	-Kaposi sarcoma -Ovarian germ cell tumor o -Testicular germ cell tumor.	
		injection: 3 mg/ml in 10-ml ampoule tablet: 15 mg -early stage colon cancer -early stage rectal cancer -early stage breast cancer -gestational trophoblastic neoplasia -metastatic colorectal cancer o-osteosarcoma o-Burkitt lymphoma	T
	calcium folinate		

Therapeutic Category	Generic Name	Formulations	Service Level Category
	capecitabine	tablet: 150 mg, 500 mg o-early stage colon cancer o-early stage rectal cancer o-metastatic breast cancer -metastatic colorectal cancer	T
	carboplatin	injection: 50 mg/5 mL, 150 mg/15 mL, 450 mg/45 mL, 600 mg/60 mL -early stage breast cancer -epithelial ovarian cancer -nasopharyngeal cancer, non-small cell lung cancer -osteosarcoma -retinoblastoma.	T
	chlorambucil	tablet: 2 mg -chronic lymphocytic leukemia.	T
	cisplatin	injection: 50 mg/50 mL, 100 mg/100 mL -cervical cancer (as a radiosensitizer) -head and neck cancer (as a radiosensitizer) -nasopharyngeal cancer (as a radio sensitizer) -non-small cell lung cancer -osteosarcoma -ovarian germ cell tumor -testicular germ cell tumor	T
	cytotoxic	powder for injection: 500 mg in vial tablet: 25 mg - chronic lymphocytic leukemia - diffuse large b-cell lymphoma - early stage breast cancer - gestational trophoblastic neoplasia - hodgkin lymphoma - follicular lymphoma - rhabdomyosarcoma - ewing sarcoma -acute lymphoblastic leukemia	T

Therapeutic Category	Generic Name	Formulations	Service Level Category
		<ul style="list-style-type: none"> - burkitt lymphoma - metastatic breast cancer. 	
		Powder for injection: 100 mg in vial	T
		<ul style="list-style-type: none"> o -acute myelogenous leukemia o- acute lymphoblastic leukemia o- acute promyelocytic leukemia 	
		o- burkitt lymphoma.	
	cytarabine	<ul style="list-style-type: none"> Powder for injection: 100 mg in vial o -hodgkin lymphoma. 	T
	dacarbazine	<ul style="list-style-type: none"> Powder for injection: 500 micrograms in vial - gestational trophoblastic neoplasia - rhabdomyosarcoma - wilms tumor. 	T
	dactinomycin	<ul style="list-style-type: none"> Powder for injection: 50 mg (hydrochloride) in vial - acute lymphoblastic leukemia - acute myelogenous leukemia - acute promyelocytic leukemia. 	T
	daunorubicin	<ul style="list-style-type: none"> injection: 20 ng/ml, 40 mg/m² o -early stage breast cancer o -metastatic breast cancer o -metastatic prostate cancer. 	T
	docetaxel	<ul style="list-style-type: none"> Powder for injection: 10 mg, 50 mg (hydrochloride) in vial o -diffuse large b-cell lymphoma o -early stage breast cancer o -hodgkin lymphoma o -kaposi sarcoma. - follicular lymphoma o -metastatic breast cancer - osteosarcoma o -ewing sarcoma 	T
	doxorubicin		

Therapeutic Category	Generic Name	Formulations	Service Level Category
		<ul style="list-style-type: none"> o -acute lymphoblastic leukemia -wilms tumor 	
		<ul style="list-style-type: none"> o -burkitt lymphoma <p>capsule: 100 mg, injection: 20 mg/ml in 5-ml ampoule:</p> <ul style="list-style-type: none"> -testicular germ cell tumor o -gestational trophoblastic neoplasia o -hodgkin lymphoma o -non-small cell lung cancer o -ovarian germ cell tumor o -retinoblastoma o -ewing sarcoma o -acute lymphoblastic leukemia o -burkitt lymphoma. 	T
etoposide		<p>powder for injection: 50 mg (phosphate) in vial tablet: 10 mg</p> <ul style="list-style-type: none"> o -chronic lymphocytic leukemia. 	T
	fludarabine	<p>injection: 50 mg/ml in 5-ml ampoule</p> <ul style="list-style-type: none"> o -early stage breast cancer o -early stage colon cancer o -early stage rectal cancer o -metastatic colorectal cancer -nasopharyngeal cancer. 	T
	fluorouracil (D)	<p>injection: 120 micrograms/0.2 ml; 300 micrograms/0.5 ml; 480 micrograms/0.8 ml in pre-filled syringe; 300 micrograms/ml in 1-ml vial; 480 mg/1.6 ml in 1.6-ml vial.</p> <ul style="list-style-type: none"> - as primary prophylaxis in patients at high risk for developing febrile neutropenia associated with myelotoxic chemotherapy. - as secondary prophylaxis for patients who experienced neutropenia following prior myelotoxic chemotherapy. 	T
	filgrastim	<ul style="list-style-type: none"> -to facilitate administration of dose-dense chemotherapy regimens. 	

Therapeutic Category	Generic Name	Formulations	Service Level Category
	gemcitabine	powder for injection: 200 mg in vial, 1 g in vial o -epithelial ovarian cancer o -non-small cell lung cancer.	T
	hydroxycarbamide	solid oral dosage form: 200 mg, 250 mg, 300 mg, 400 mg, 500 mg, 1 gram o -chronic myeloid leukemia. powder for injection: 500 mg vial, 1-gram vial, 2-gram vial o-ovarian and testicular germ cell tumor	T
	ifosfamide	o-osteosarcoma o -rhabdomyosarcoma o-ewing sarcoma.	
	imatinib	tablet: 100 mg, 400 mg -chronic myeloid leukemia -gastrointestinal stromal tumor.	T
	irinotecan	injection: 40 mg/2 ml in 2-ml vial; 100 mg/5 ml in 5-ml vial; 500 mg/25 ml in a 25-ml vial o -metastatic colorectal cancer.	T
	mercaptopurine	tablet: 50 mg o -acute lymphoblastic leukemia o -acute promyelocytic leukemia.	T
	mesna	injection: 100 mg/ml in 4-ml and 10-ml ampoules tablet: 400 mg, 600 mg o -testicular germ cell tumor o -ovarian germ cell tumor o -osteosarcoma o -rhabdomyosarcoma o-ewing sarcoma.	T
	methotrexate (D)	powder for injection: 50 mg (as sodium salt) in vial tablet: 2.5 mg (as sodium salt) o -early stage breast cancer o -gestational trophoblastic neoplasia -osteosarcoma	T

Therapeutic Category	Generic Name	Formulations	Service Level Category
		-acute lymphoblastic leukemia o- acute promyelocytic leukemia. injection: 50 mg/10 ml in 10-ml vial; 100 mg/20 ml in 20-ml vial, 200 mg/40 ml in 40-ml vial	T
	oxaliplatin	o-early stage colon cancer o-metastatic colorectal cancer. powder for injection: 6 mg/ml - epithelial ovarian cancer	T
		o-early stage breast cancer o-metastatic breast cancer o-kaposi sarcoma	
	paclitaxel	o-nasopharyngeal cancer o-non-small cell lung cancer -ovarian germ cell tumor.	T
		capsule: 50 mg (as hydrochloride) injection: 100 mg/10 ml in 10-ml vial; 500 mg/50 ml in 50-ml vial -diffuse large b-cell lymphoma	T
	rituximab	o- chronic lymphocytic leukemia o-follicular lymphoma.	T
	thioguanine [c]	solid oral dosage form: 40 mg o-acute lymphoblastic leukemia.	T
	trastuzumab	powder for injection: 60 mg; 40 mg in vial o-early stage HER2 positive breast cancer o-metastatic HER2 positive breast cancer.	T
	vinblastine	powder for injection: 10 mg (sulfate) in vial o-hodgkin lymphoma o-kaposi sarcoma o-testicular germ cell tumor. -ovarian germ cell tumor.	T
	vincristine	powder for injection: 1 mg, 5 mg (sulfate) in vial o-diffuse large b-cell lymphoma	T

Therapeutic Category	Generic Name	Formulations	Service Level Category
		<ul style="list-style-type: none"> o -gestational trophoblastic neoplasia o -hodgkin lymphoma o -Kaposi sarcoma o -follicular lymphoma o -retinoblastoma o -rhabdomyosarcoma, o -ewing sarcoma o -acute lymphoblastic leukemia o -wilms tumor o -burkitt lymphoma. 	
	vinorelbine	<ul style="list-style-type: none"> injection: 10 mg/ml in 1-ml vial; 50 mg/5 ml in 5-ml vial - non-small cell lung cancer. - metastatic breast cancer 	T
8.3. Hormones and antihormones			
	<input type="checkbox"/> anastrozole	<ul style="list-style-type: none"> tablet: 1 mg - early stage breast cancer - metastatic breast cancer 	T
	<input type="checkbox"/> bicalutamide	<ul style="list-style-type: none"> tablet: 50 mg - metastatic prostate cancer 	T
		<ul style="list-style-type: none"> injection: 4 mg/ml in 1-ml ampoule (as disodium phosphate salt) oral liquid: 2 mg/5 ml [C] -acute lymphoblastic leukemia 	P, S, T
	<input type="checkbox"/> dexamethasone (D)	Dose form:	
	<input type="checkbox"/> leuprorelin	<ul style="list-style-type: none"> - early stage breast cancer - metastatic prostate cancer 	T
	hydrocortisone (D)	<ul style="list-style-type: none"> Powder for injection: 100 mg (as sodium succinate) in vial -acute lymphoblastic leukemia 	P, S, T
	methylprednisolone [C]	<ul style="list-style-type: none"> Injection: 40 mg/mL (as sodium succinate) in 1-mL single-dose vial and 5-mL multi-dose vials; 80 mg/mL (as sodium succinate) in 1-mL single-dose vial. o - acute lymphoblastic leukemia. 	S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
		oral liquid: 5 mg/ml [C] tablet: 5 mg, 25 mg - chronic lymphocytic leukemia - diffuse large b-cell lymphoma - hodgkin lymphoma - follicular lymphoma - acute lymphoblastic leukemia - burkitt lymphoma.	S, T
	<input type="checkbox"/> prednisolone (D)	tablet: 10 mg, 20 mg (as citrate) - early stage breast cancer - metastatic breast cancer.	T
	tamoxifen		
9. Anti Parkinsonism Medicines		injection: 5 mg (lactate) in 1-ml ampoule tablet: 2 mg (hydrochloride)	S, T S, T
10. Medicines Affecting the Blood			
10.1. Antianaemia medicines			
	ferrous salt	oral liquid: equivalent to 25 mg iron (as sulfate)/ml tablet: equivalent to 60 mg iron	P, S, T P, S, T
	ferrous salt + folic acid	tablet: equivalent to 60 mg iron + 400 micrograms folic acid (nutritional supplement for use during pregnancy)	P, S, T
	folic acid*	tablet: 400 micrograms*, 1 mg, 5 mg. *use for periconceptual to prevent the first occurrence of neural tube defects.	P, S, T
	hydroxocobalamin	injection: 1 mg (as acetate, as hydrochloride or as sulfate) in 1-ml ampoule	P, S, T
	tirofiban (P)	injection: 0.25 mg/ml.	T
	iron sucrose (P)	injection: 20 mg/ml.	S, T
10.2. Medicines affecting coagulation			
	<input type="checkbox"/> enoxaparin*	injection: ampoule or pre-filled syringe 20 mg/0.2 ml; 40 mg/0.4 ml; 60 mg/0.6 ml; 80 mg/0.8 ml; 100 mg/1 ml; 120 mg/0.8 ml; 150 mg/1 ml	T
	clopidogrel	*alternatives are limited to nadroparin and dalteparin. tablet: 75 mg	S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
	heparin sodium (D)	injection: 1000 IU/ml; 5000 IU/ml; 20 000 IU/ml in 1-ml ampoule injection: 1 mg/ml ampoule; [E] 10 mg/ml in 5-ml tablet: 10 mg	S, T
	phycomenadione	tablet: 10 mg	P, S, T
	protamine sulfate (D)	injection: 10 mg/ml in 5-ml ampoule	P, S, T
	tranexamic acid	injection: 100 mg/ml in 10-ml ampoule capsules: 250 mg, 500 mg	S, T
	□ warfarin (D)	tablet: 1 mg, 2 mg, 5 mg (sodium salt) injection: 4 micrograms/ml (as acetate) in 1-ml ampoule	P, S, T
	desmopressin	nasal spray: 10 micrograms (as acetate) per dose	S, T
10.3. Other medicines for haemoglobinopathies			
	deferoxamine*	powder for injection: 500 mg (mesilate) in vial *deferasirox oral form may be an alternative, depending on cost and availability.	S, T
	hydroxycarbamide	solid oral dosage form: 200 mg, 500 mg, 1 gram	S, T
II. Blood Products of Human Origin and Plasma Substitutes			
II.1. Blood and blood components			
In accordance with the World Health Assembly resolution (WHA63.12), WHO recognizes that achieving self-sufficiency—unless special circumstances prevent it—in the supply of safe blood components, based on voluntary, non-remunerated blood donation; and the security of that supply, are important national goals to prevent blood shortages and to meet the transfusion requirements for the patient population. All preparations should comply with the WHO requirements.			
	fresh-frozen plasma		T
	platelets		T
	red blood cells		S, T
	whole blood		T
II.2. Plasma-derived medicines			
All human plasma-derived medicines should comply with WHO requirements			
II.2.1 Human immunoglobulins		injection: 250 micrograms in a single-dose vial	S, T
	anti-D immunoglobulin	injection: 150 IU/ml in vial	S, T
	anti-rabies immunoglobulin	injection: 500 IU in vial	S, T
	anti-tetanus immunoglobulin	intramuscular administration: 16% protein solution	S, T
	normal immunoglobulin	* indicated for primary immune deficiency. intravenous administration: 5%, 10% protein solution	T

Therapeutic Category	Generic Name	Formulations	Service Level Category
		*indicated for primary immune deficiency and kawasaki disease subcutaneous administration: 15%, 16% protein solution	T
		* indicated for primary immune deficiency. powder for injection: 500 IU/vial	S, T
		powder for injection: 500 IU/vial, 1000 IU/vial	S, T
11.3. Plasma substitutes			
	<input type="checkbox"/> dextran 70*	injectable solution: 6%	P, S, T
	<input type="checkbox"/> dextran 40 (P)	*bolivene, injectable solution; 3.5% is considered as equivalent. injection: 10%.	S, T
12. Cardiovascular Medicines			
12.1. Antianginal medicines			
	<input type="checkbox"/> atenolol (D)	tablet: 50 mg and 100 mg.	P, S, T
	<input type="checkbox"/> glyceryl trinitrate	tablet (sublingual): 500 micrograms.	P, S, T
	<input type="checkbox"/> isosorbide dinitrate	tablet (sublingual): 5 mg.	P, S, T
	<input type="checkbox"/> losartan potassium	tablet: 50mg	S
	<input type="checkbox"/> propranolol	tablets: 10mg, 40mg, 80 mg	S
	<input type="checkbox"/> verapamil (D)	tablet: 40 mg, 80 mg (hydrochloride).	S, T
12.2. Antiarrhythmic medicines			
	<input type="checkbox"/> bisoprolol (D)	tablet: 1.25 mg, 5 mg. * includes metoprolol and carvedilol as alternatives.	P, S, T
	<input type="checkbox"/> digoxin (D)	injection: 250 micrograms/ml in 2-ml ampoule. oral liquid: 50 micrograms/ml.	S, T
	<input type="checkbox"/> epinephrine (adrenaline) (D)	tablet: 62.5 micrograms, 250 micrograms. injection: 100 micrograms/ml (as acid tartrate or hydrochloride) in 10-ml ampoule.	S, T
	<input type="checkbox"/> lidocaine	injection: 20 mg (hydrochloride)/ml in 5-ml ampoule.	P, S, T
	<input type="checkbox"/> verapamil (D)	injection: 2.5 mg (hydrochloride)/ml in 2-ml ampoule. tablet: 40 mg, 80 mg (hydrochloride).	S, T
	<input type="checkbox"/> amiodarone	injection: 50 mg/ml in 3-ml ampoule (hydrochloride). tablet: 100 mg, 200 mg, 400 mg (hydrochloride).	S, T
	<input type="checkbox"/> adenosine (P)	injection: 3 mg/ml	S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
12.3. Antihypertensive medicines		intravenous injection: 5 mg/ml.	P, S, T
	<input type="checkbox"/> amlodipine	tablet: 5 mg (as maleate, mesylate, or besylate). tablet: 1.25 mg, 5 mg. *includes atenolol, metoprolol, and carvedilol as alternatives. do not use atenolol as a first line agent in uncomplicated hypertension in patients >60 years.	P, S, T
	<input type="checkbox"/> bisoprolol (D)*	tablets: 10 mg, 40 mg, 80 mg. Injection: 20 mg tablet: 2.5 mg, 5 mg (as hydrogen maleate). powder for injection: 20 mg (hydrochloride) in ampoule. tablet: 25 mg, 50 mg (hydrochloride).	P, S, T
	<input type="checkbox"/> enalapril (D)	*Use hydralazine only for acute management of severe pregnancy-induced hypertension. Using it to treat essential hypertension is not recommended because other medicines have shown evidence of greater efficacy and safety.	P, S, T
	<input type="checkbox"/> hydralazine*	oral liquid: 50 mg/5 ml. solid oral dosage form: 12.5 mg, 25 mg. injection: 20 mg. tablet: 250 mg. injection: 250 mg.	S, T
	<input type="checkbox"/> hydrochlorothiazide(D)	*Use methyl/dopa only to manage pregnancy-induced hypertension. Using it to treat essential hypertension is not recommended because other medicines have shown greater efficacy and safety.	P, S, T
	<input type="checkbox"/> isoprenaline (P)	powder for infusion: 50 mg in ampoule. injection 1mg/ml (iv):	T
12.4. Medicines used in heart failure			
	<input type="checkbox"/> bisoprolol* (D)	tablet: 1.25 mg, 5 mg. *includes metoprolol and carvedilol as alternatives.	P, S, T
	<input type="checkbox"/> digoxin (D)	injection: 250 micrograms/ml in 2-ml ampoule. oral liquid: 250 micrograms/ml.	S, T
	<input type="checkbox"/> enalapril (D)	tablet: 62.5 micrograms, 250 micrograms. tablet: 2.5 mg, 5 mg (as hydrogen maleate).	S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
		injection: 10 mg/ml in 2-ml ampoule. oral liquid: 20 mg/5 ml [c]. tablet: 40 mg.	S, T
	<input type="checkbox"/> furosemide (D)	oral liquid: 50 mg/5 ml. solid oral dosage form: 25 mg.	P, S, T
	<input type="checkbox"/> hydrochlorothiazide (D)	tablet: 25 mg.	P, S, T
	spironolactone	injection: 40 mg/ml(<i>hydrochloride</i>) in 5-ml vial	P, S, T
	dopamine	250 mg injection.	S, T
	dobutamine	injection: 30 mg/ml.	S, T
	ephedrine (P)		S, T
12.5. Antithrombotic medicines			
12.5.1 Anti-platelet medicines	aceylsalicylic acid clopidogrel	tablet: 100 mg. tablet: 75 mg, 300 mg.	P, S, T
12.5.2 Thrombolytic medicines	streptokinase	powder for injection: 1.5 million IU in vial.	T
12.6. Lipid-lowering agents			
	<input type="checkbox"/> simvastatin*	tablet: 5 mg, 10 mg, 20 mg, 40 mg. *use in high-risk patients.	T
	atorvastatin (P)	tablet: 10 mg, 20 mg, 40 mg.	S, T
	risovastatin (P)	tablet: 10 mg, 20 mg, 40 mg.	S, T
13. Dermatological Medicines (topical)			
13.1. Antifungal medicines			
	<input type="checkbox"/> miconazole	cream or ointment: 2% (<i>nitrate</i>).	P, S, T
	selenium sulfide	detergent-based suspension: 2%.	P, S, T
	sodium thiosulfate	solution: 15%.	P, S, T
	terbinafine	cream: 1% or ointment: 1% terbinafine (<i>hydrochloride</i>).	P, S, T
13.2. Anti-infective medicines			
	mupirocin	cream (as mupirocin calcium): 2%. ointment: 2%.	P, S, T
	potassium permanganate	aqueous solution: 1:10 000.	P, S, T
	silver sulfadiazine (a)	cream: 1%.	P, S, T

Therapeutic Category	Generic Name	(a) >2 months.	Formulations	Service Level Category
13.3. Anti-inflammatory and antipruritic medicines				
	<input type="checkbox"/> betamethasone (a)	cream or ointment: 0.1% (as valerate). a hydrocortisone is preferred for neonates.	P, S, T	
	<input type="checkbox"/> calamine	lotion.	P, S, T	
	<input type="checkbox"/> hydrocortisone	cream or ointment: 1% (acetate).	P, S, T	
13.4. Medicines affecting skin differentiation and proliferation				
	benzoyl peroxide	cream or lotion: 5%.	P, S, T	
	coal tar	solution: 5%.	P, S, T	
	fluorouracil (D)	ointment: 5%.	S, T	
	<input type="checkbox"/> podophyllin resin	solution: 10% to 25%.	S, T	
	<input type="checkbox"/> salicylic acid	solution: 5%.	P, S, T	
	salicylic acid + betamethasone	lotion: 3% + 0.5 % w/v	S, T	
	urea	cream or ointment: 5%, 10%.	P, S, T	
13.5. Scabicides and pediculicides				
	<input type="checkbox"/> benzyl benzoate (a)	lotion: 25%. (a) >2 years.	P, S, T	
	permethrin	cream: 5%.	P, S, T	
		lotion: 1%.	P, S, T	
13.6. Skin treatment				
	silver sulfadiazine	cream: 1%.	P, S, T	
	sodium thiosulfate	solution: 15%.	P, S, T	
14. Diagnostic Agents				
14.1. Ophthalmic medicines				
	fluorescein	eye drops: 1% (sodium salt).	S, T	
	<input type="checkbox"/> tropicamide	eye drops: 0.5%.	S, T	
14.2. Radio-contrast media				
	<input type="checkbox"/> amidotriozate	injection: 140 mg to 420 mg iodine (as sodium or meglumine salt)/ml in 20-ml ampoule.	S, T	
	barium sulfate	aqueous suspension.	S, T	
	<input type="checkbox"/> iohexol	injection: 140 mg to 350 mg iodine/ml in 5-ml, 10ml, 20-ml ampoules.	S, T	
	barium sulfate [c]	aqueous suspension	S, T	

Therapeutic Category	Generic Name	Formulations	Service Level Category
15. Disinfectants and Antiseptics		solution: 5 grams to 8 grams iodine in 100 ml to 250 ml.	S, T
15.1. Antiseptics			
	<input type="checkbox"/> chlorhexidine	solution: 5% digluconate.	P, S, T
	<input type="checkbox"/> ethanol	solution: 70% (denatured).	P, S, T
	<input type="checkbox"/> povidone iodine	solution: 10% (equivalent to 1% available iodine).	P, S, T
15.2. Disinfectants			
		solution containing ethanol 80% volume /volume, solution containing isopropyl alcohol 75% volume/volume.	P, S, T
		powder: (0.1% available chlorine) for solution.	P, S, T
		solution: 4.8%.	P, S, T
		solution: 2%.	P, S, T
		solution	P, S, T
16. Diuretics			
		alcohol-based hand rub	S, T
		<input type="checkbox"/> chlorine base compound	P, S, T
		<input type="checkbox"/> chloroxylenol	P, S, T
		glutaral	P, S, T
		methylated spirit	P, S, T
		amiloride	S, T
		<input type="checkbox"/> furosemide (D)	P, S, T
		hydrochlorothiazide (D)	P, S, T
		<input type="checkbox"/> hydrochlorothiazide (D)	P, S, T
		mannitol (D)	P, S, T
		spironolactone	P, S, T
		sodium Polystyrene sulfonate (P)	P, S, T
17. Gastrointestinal Medicines			
17.1. Antidiarrhoeals		Age-appropriate formulations and doses including lipase, protease, and amylase.	T
		<input type="checkbox"/> pancreatic enzymes	
		<input type="checkbox"/> omeprazole	P, S, T
		<input type="checkbox"/> ranitidine	P, S, T
		powder for injection: 40 mg in vial.	P, S, T
		powder for oral liquid: 20 mg, 40 mg sachets	P, S, T
		solid oral dosage form: 10 mg, 20 mg, 40 mg.	P, S, T
		injection: 25 mg/ml (as hydrochloride) in 2-ml ampoule.	P, S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
		oral liquid: 75 mg/5 ml (as hydrochloride). tablet: 150 mg (as hydrochloride).	P, S, T
			P, S, T
17.2. Antiemetic medicines			
	dexamethasone (D)	injection: 4 mg/ml in 1-ml ampoule (as disodium phosphate salt). oral liquid: 0.5 mg/5 ml, 2 mg/5 ml. solid oral dosage form: 0.5 mg, 0.75 mg, 1.5 mg, 4 mg.	P, S, T
		injection: 5 mg (hydrochloride)/ml in 2-ml ampoule. oral liquid: 5 mg/5 ml [C]. tablet: 10 mg (hydrochloride).	P, S, T
	metoclopramide (a)	(a) Do not use in neonates. injection: 2 mg base/ml in 2-ml ampoule (as hydrochloride). oral liquid: 4 mg base/5 ml. solid oral dosage form: equivalent to 4 mg base, equivalent to 8 mg base, equivalent to 24 mg base. (a) > 1 month.	P, S, T
	ondansetron (a)		S, T
			S, T
			S, T
17.3. Anti-inflammatory medicines			
	<input type="checkbox"/> sulfasalazine (D)	retention enema. suppository: 500 mg. tablet: 500 mg.	S, T
	<input type="checkbox"/> hydrocortisone	retention enema. suppository: 25 mg (acetate). (the <input type="checkbox"/> only applies to hydrocortisone retention enema.)	P, S, T
17.4. Laxatives			S, T
	<input type="checkbox"/> senna	tablet: 7.5 mg (sennosides) (or traditional dosage forms).	P, S, T
17.5. Medicines used in diarrhea			
	oral rehydration salts (D)	dry mixture (low osmolarity formula) in sachet for 1 liter of solution; each sachet contains glucose anhydrous 13.5gm B.P., trisodium citrate dihydrate 2.9 gm B.P., potassium chloride 1.5gm B.P., sodium chloride 2.6gm B.P.	P, S, T
	zinc sulfate [*]	solid oral dosage form: 20 mg.	P, S, T
		*For acute diarrhea, use zinc sulfate as an adjunct to oral rehydration salts.	

Therapeutic Category	Generic Name	Formulations	Service Level Category
17.6. Other Medicines 16			
	aluminum hydroxide + magnesium trisilicate	suspension: 215mg + 80mg / 5ml tablet: 5 mg	P, S, T
	bisacodyl	injection: 40mg tablet: 50 mg	P, S, T
	dimenhydrinate		P, S, T
	glycerin	suppository	P, S, T
	magnesium oxides and hydroxides	suspension: 7.9 %w/v	P, S, T
	phloroglucinol	Injection: 10mg /ml	P, S, T
	sodium biphosphate	Enema: 7.2g /120ml	P, S, T
18. Hormones, other Endocrine Medicines and Contraceptives			
18.1. Adrenal hormones and synthetic substitutes			
	fluocortisone	tablet: 100 micrograms (acetate),	T
	hydrocortisone (D)	tablet: 5 mg, 10 mg, 20 mg.	S, T
18.2. Androgens			
	testosterone	injection: 200 mg (enanthate) in 1-ml ampoule.	S, T
18.3. Contraceptives			
	<input type="checkbox"/> ethynodiol + <input type="checkbox"/> levonorgestrel	tablet: 30 micrograms + 150 micrograms.	P, S, T
	<input type="checkbox"/> ethynodiol + <input type="checkbox"/> norethisterone	tablet: 35 micrograms + 1 mg	P, S, T
	levonorgestrel (D)	tablet: 30 micrograms, 750 micrograms (pack of 2), 1.5 mg.	P, S, T
	estradiol cypionate + medroxyprogesterone acetate (D)	injection: 5 mg + 25 mg. depot injection: 150 mg/ml in 1-ml vial.	S, T
	norethisterone enantate	oily solution: 200 mg/ml in 1-ml ampoule.	S, T
18.3.3. Copper-containing device			P, S, T
18.3.3.1. Intrauterine devices		levonorgestrel-releasing intrauterine system with reservoir with 52 mg of levonorgestrel.	T
18.3.4 Barrier methods	condoms diaphragm		P, S, T
	etonogestrel-releasing implant	single-rod etonogestrel-releasing implant with 68 mg of etonogestrel.	S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
18.3.5 Implantable contraceptives	levonorgestrel-releasing implant (D)	two-rod levonorgestrel-releasing implant; each rod contains 75 mg of levonorgestrel (150 mg total).	S, T
18.3.6 Intravaginal contraceptive	progesterone vaginal ring*	progesterone-releasing vaginal ring with 2.074 grams of micronized progesterone. *use for women actively breastfeeding at least 4 times per day.	S, T
18.4. Estrogen	estrogen		S, T
18.5. Insulin and other medicines used for diabetes			
	<input type="checkbox"/> gliclazide*	solid oral dosage form: (controlled release tablets) 30 mg, 60 mg, 80 mg. *glibenclamide not suitable above 60 years.	S, T
	<input type="checkbox"/> glucagon	injection: 1 mg/ml.	T
	insulin injection (soluble)	injection: 40 iu/ml in 10-ml vial, 100 iu/ml in 10-ml vial.	P, S, T
	intermediate-acting insulin	injection: 40 iu/ml in 10-ml vial, 100 iu/ml in 10-ml vial (as compound insulin zinc suspension or isophane insulin).	P, S, T
	metformin	tablet: 500 mg (hydrochloride).	P, S, T
	glibenclamide	tablets 5 mg	P, S, T
18.6. Ovulation inducer			
	clomifene	tablet: 50 mg (citrate).	S, T
18.7. Progestogen	<input type="checkbox"/> medroxyprogesterone	tablet: 5 mg acetate.	P, S, T
18.8. Thyroid hormone and antithyroid medicine			
	levothyroxine	tablet: 25 micrograms [c], 50 micrograms, 100 micrograms (sodium salt)	P, S, T
	potassium iodide	tablet: 60 mg.	P, S, T
	<input type="checkbox"/> propylthiouracil	tablet: 50 mg.	S, T
	Lugol's solution	oral liquid: about 130 mg total iodine/ml.	P, S, T
	propylthiouracil	tablet: 50 mg.	P, S, T
19. Immunologicals			
19.1. Diagnostic agents	tuberculin, purified protein derivative (PPD)	injection	P, S, T
		All tuberculins should comply with the WHO requirements for tuberculins.	

Therapeutic Category	Generic Name	Formulations	Service Level Category
19.2.	Sera and immunoglobulins		
All plasma fractions should comply with the WHO requirements.	anti-D immunoglobulin (human) anti-venom immunoglobulin* anti-hepatitis B immunoglobulin anti-rabies immunoglobulin (human) anti-tetanus immunoglobulin (human) diphtheria antitoxin	injection: 250 microgram in single-dose vial. injection: *exact type to be defined locally. injection injection: 150 IU/ml in vial. injection: 500 IU in vial. injection: 10 000 IU, 20 000 IU in vial.	P, S, T P, S, T P, S, T P, S, T P, S, T S, T
19.3.	Vaccines		
All vaccines under the Expanded Program on Immunization (EPI) of Pakistan will be included in the NEML. All vaccines must comply with the WHO requirements for biological substances and should be administered under WHO guidelines. WHO stated that vaccines used in children should be polyvalent. The new vaccines, which the National Technical Advisory Group on Immunization (NTACI) approved, and planned to be given under the EPI, will be included in the NEML as and when listed in the EPI. In the future, the vaccines under consideration, if and when they are included in the EPI, will also be included in the NEML, from the date of inclusion.	BCG vaccine diphtheria vaccine Haemophilus influenzae type b vaccine hepatitis B vaccine HPV vaccine measles vaccine pertussis vaccine pneumococcal vaccine poliomyelitis vaccine rotavirus vaccine rubella vaccine tetanus vaccine tetanus immunoglobulin (human) meningococcal meningitis vaccine yellow fever vaccine cholera vaccine hepatitis A vaccine meningococcal meningitis vaccine	Recommendations for all	P, S, T S, T Recommendations for certain regions Recommendations for some high-risk populations

Therapeutic Category	Generic Name	Formulations	Service Level Category
	rabies vaccine	vial: 150 IU/ml	
	typhoid vaccine		
	influenza vaccine (seasonal)		Recommendations for immunization programs with certain characteristics
	mumps vaccine		
	varicella vaccine		
20. Muscle Relaxants			
	<input type="checkbox"/> atracurium	injection: 10 mg/ml (besylate). injection: 500 micrograms in 1-ml ampoule, 2.5 mg (metilsulfate) in 1-ml ampoule.	S, T
	<input type="checkbox"/> neostigmine	tablet: 15 mg (bromide).	S, T
	<input type="checkbox"/> suxamethonium	injection: 50 mg (chloride)/ml in 2-ml ampoule. powder for injection (chloride) in vial.	S, T
	<input type="checkbox"/> vecuronium [c]	powder for injection: 10 mg (bromide) in vial.	S, T
	<input type="checkbox"/> pyridostigmine	injection: 1 mg in 1-ml ampoule, tablet: 60 mg (bromide).	P, S, T
	<input type="checkbox"/> pancuronium	injection: contains pancuronium bromide 2mg/ml, 2 ml.	S, T
21. Ophthalmological Preparations			
21.1. Anti-infective agents			
	<input type="checkbox"/> aciclovir (D)	ointment: 3% w/v. eye drop 4% w/v.	S, T
	<input type="checkbox"/> sodium cromoglycate		S, T
	<input type="checkbox"/> gentamicin (D)	solution (eye drops): 0.3% (sulfate).	P, S, T
	<input type="checkbox"/> ofloxacin	solution (eye drops): 0.3%.	P, S, T
	<input type="checkbox"/> tetracycline	eye ointment: 1% (hydrochloride).	P, S, T
	<input type="checkbox"/> azithromycin	solution:1.50%	T
	<input type="checkbox"/> erythromycin*	ointment:0.50%	
	<input type="checkbox"/> natamycin	suspension:5%	T
	<input type="checkbox"/> ciprofloxacin	eye drops: 0.3 %w/v	T
	<input type="checkbox"/> dexamethasone	eye drops: 0.1 %w/v	T
	<input type="checkbox"/> miconazole (nitrate)	eye ointment:0.02	T

Therapeutic Category	Generic Name	Formulations	Service Level Category
	polymyxin B (sulphate) + bacitracin zinc	eye ointment: 10000IU/g + 500IU/g	T
	proparacaine (hydrochloride)	drops: 0.005	T
	tears natural	eye drops	T
	tobramycin + dexamethasone	eye drops: 0.3 % + 0.1% w/v	T
	tropicamide	eye drops: 1 %w/v	T
	timolol (hydrogen maleate)	eye drops: 0.25%; 0.5%	T
21.2. Anti-inflammatory agents			
	<input type="checkbox"/> prednisolone	solution (eye drops): 0.5% (sodium phosphate).	P, S, T
21.3. Local anesthetics			
	<input type="checkbox"/> tetracaine (a)	solution (eye drops): 0.5% (hydrochloride). (a) not in preterm neonates.	P, S, T
21.4. Miotics and antiglaucoma medicines			
	acetazolamide	tablet: 250 mg.	S, T
	latanoprost	solution (eye drops): latanoprost 50 micrograms/ml.	S, T
	<input type="checkbox"/> pilocarpine	solution (eye drops): 2%, 4% (hydrochloride or nitrate).	S, T
	<input type="checkbox"/> timolol	solution (eye drops): 0.25%, 0.5% (as hydrogen maleate).	P, S, T
21.5. Mydriatics			
	atropine (a)*	solution (eye drops): 0.1%, 0.5%, 1% (sulfate). *[c] or homatropine (hydrobromide), or cyclopentolate (hydrochloride); (a) >3 months.	P, S, T
	epinephrine (adrenaline) (D)	Solution (eye drops): 2% (as hydrochloride).	P, S, T
21.6. Antivascular endothelial growth factor (VEGF) preparations			
	bevacizumab	injection: 25 mg/ml.	T
22. Oxytocics and Anti-Oxytocics			
22.1. Oxytocics			
	<input type="checkbox"/> ergometrine	injection: 200 micrograms (hydrogen maleate) in 1ml ampoule. tablet: 200 micrograms.	P, S, T
	misoprostol	- to manage incomplete abortion and miscarriage. - to prevent and treat postpartum hemorrhage if oxytocin is not available or cannot be safely used vaginal tablet: 25 micrograms *	S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
		*only use to induce labor if appropriate facilities are available.	
	oxytocin	injection: 10 IU in 1-ml.	S, T
22.2. Antioxytoxics (tocolytics)	nifedipine	capsule: immediate release, 10 mg. tablet: slow release, 20 mg.	S, T T
23. Peritoneal Dialysis Solution	intraperitoneal dialysis solution (of appropriate composition).	parenteral solution.	S, T
24. Medicines for Mental and Behavioral Disorder			
24.1. Medicines used in psychotic disorders			
	<input type="checkbox"/> chlorpromazine (D)	injection: 25 mg (hydrochloride)/ml in 2-ml ampoule. oral liquid: 25 mg (hydrochloride)/5 ml.	P, S, T
	<input type="checkbox"/> fluphenazine	tablet: 10 mg, 25 mg, 50 mg, 100 mg (hydrochloride).	P, S, T
	<input type="checkbox"/> haloperidol (D)	injection: 25 mg (decanoate or enanthate) in 1-ml ampoule. injection: 5 mg in 1-ml ampoule. oral liquid: 2 mg/ml.	S, T
	<input type="checkbox"/> risperidone	tablet: 0.5 mg, 2 mg, 5 mg.	S, T
	<input type="checkbox"/> clozapine	solid oral dosage form: 0.25 mg to 6.0 mg. solid oral dosage form: 25 to 200 mg.	S, T
24.2. Medicines used in mood disorders			
24.2.1 Medicines used in depressive disorders	<input type="checkbox"/> amitriptyline	tablet: amitriptyline hcl 10 mg, 25 mg. solid oral dosage form: 20 mg (as hydrochloride).	P, S, T
	<input type="checkbox"/> fluoxetine (D) (a)	(a) >8 years	P, S, T
24.2.2 Medicines used in bipolar disorders	carbamazepine (D)	tablet (scored): 100 mg, 200 mg.	P, S, T
	lithium carbonate	syrup: 100 mg/5 ml.	P, S, T
	valproic acid (sodium valproate) (D)	solid oral dosage form: 300 mg. tablet (enteric-coated): 200 mg, 500 mg (sodium valproate).	T
24.2.3 Others	olanzapine	tablets: 5mg, 10 mg	S, T
24.3. Medicines for anxiety disorders			S, T
	<input type="checkbox"/> diazepam	tablet (scored): 2 mg, 5 mg.	P, S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
24.4. Medicines used for obsessive compulsive disorders			
	<input type="checkbox"/> clomipramine	capsule: 10 mg, 25 mg (hydrochloride).	P, S, T
24.5. Medicines for disorders due to psychoactive substance use			
		chewing gum: 2 mg, 4 mg (as polacrilex).	T
		transdermal patch: 5 mg to 30 mg/16 hours, 7 mg to 21 mg/24 hours.	
		concentrate for oral liquid: 5 mg/ml, 10 mg/ml (hydrochloride).	
		oral liquid: 5 mg/5 ml, 10 mg/5 ml (hydrochloride).	T
		*the square box is added to buprenorphine. use the medicines only within an established support program.	
25. Medicines Acting on the Respiratory Tract			
25.1. Anti-asthmatic and medicines for chronic obstructive pulmonary disease			
		inhalation (aerosol): 50 micrograms (dipropionate) per dose, 100 micrograms (dipropionate) per dose, 100 micrograms (dipropionate) per dose (as cfc free forms).	S, T
		respirator solution for use in nebulizers: 800 micrograms /2 ml.	S, T
		inhalation (aerosol): 100 micrograms per dose, 200 micrograms per dose.	S, T
	<input type="checkbox"/> beclometasone	injection: 1 mg (as hydrochloride or hydrogen tartrate) in 1-ml ampoule.	P, S, T
	<input type="checkbox"/> budenoside [c] (D)	inhalation (aerosol): 20 micrograms/metered doses.	P, S, T
	<input type="checkbox"/> salbutamol	inhalation (aerosol): 100 micrograms (as sulfate) per dose.	P, S, T
		injection: 50 micrograms (as sulfate)/ml in 5-ml ampoule.	P, S, T
		metered dose inhaler (aerosol): 100 micrograms (as sulfate) per dose.	P, S, T
		respirator solution for use in nebulizers: 5 mg (as sulfate)/ml.	P, S, T
		tablets: 100 mg.	P, S, T
		sachet: 4 mg/sachet.	P, S, T
		tablet: 4 mg, 5 mg, 10 mg.	P, S, T
		injection: 25 mg/ml.	P, S, T
		tablets: 100 mg, 200 mg.	P, S, T
		syrup: 32 mg/5 ml	P, S, T

Therapeutic Category	Generic Name	Formulations	Service Level Category
26. Solution Correcting, Water, Electrolyte, and Acid Base Balance			
26.1. Oral	oral rehydration salts	powder for solution.	P, S, T
26.2. Parenteral			
glucose	injectable solution: 5% (isotonic), 10% (hypertonic), 50% (hypertonic). Injectable solution: 4% glucose, 0.18% sodium chloride (equivalent to Na+ 30 mmol/L, Cl- 30 mmol/L). Injectable solution: 5% glucose, 0.9% sodium chloride (equivalent to Na+ 150 mmol/L and Cl- 150 mmol/L), 5% glucose, 0.45% sodium chloride (equivalent to Na+ 75 mmol/L and Cl- 75 mmol/L) [Cl-]. solution: 11.2% in 20-ml ampoule (equivalent to K+ 1.5 mmol/ml, Cl- 1.5 mmol/ml).		P, S, T
glucose with sodium chloride	solution for dilution: 7.5% (equivalent to K+ 1 mmol/ml and Cl mmol/ml) [Cl-] 15% (equivalent to K2 mmol/ml and Cl2 mmol/ml) [Cl-]		P, S, T
sodium chloride (D)	injectable solution: 0.9% isotonic (equivalent to Na+ 154 mmol/l, Cl- 154 mmol/l).		P, S, T
sodium chloride	injectable solution: 1.4% isotonic (equivalent to Na+ 167 mmol/l, HCO3- 167 mmol/l). solution: 8.4% in 10-ml ampoule (equivalent to Na+ 1000 mmol/l, HCO3- 1000 mmol/l).		P, S, T
sodium hydrogen carbonate	injectable solution		P, S, T
sodium lactate, compound solution (o)	injectable solution		P, S, T
mannitol	injectable solution: 20% w/v (500 ml).		P, S, T
26.3. Miscellaneous			
	water for injection	2-ml, 5-ml, 10-ml ampoules.	
27. Vitamins and Minerals			
ascorbic acid	tablet: 50 mg.		P, S, T
calcium	tablet: 500 mg (elemental). oral liquid: 400 IU/ml.		P, S, T
cholecalciferol* [c]	solid oral dosage form: 400 IU, 1000 IU.		P, S, T

*use ergocalciferol as an alternative.

Therapeutic Category	Generic Name	Formulations	Service Level Category
	<input type="checkbox"/> ergocalciferol	oral liquid: 250 micrograms/ml (10 000 IU/ml). solid oral dosage form: 1.25 mg (50 000IU).	P, S, T
iodine		capsule: 200 mg. iodized oil: 1 ml (480 mg iodine), 0.5 ml (240 mg iodine) in ampoule (oral or injectable), 0.57 ml (308 mg iodine) in dispenser bottle. tablet: 50 mg.	P, S, T
	<input type="checkbox"/> nicotinamide	tablet: 25 mg (hydrochloride).	P, S, T
pyridoxine		capsule: 50 000 IU, 100 000 IU, 200 000 IU (as palmitate).	P, S, T
		oral oily solution: 100 000 IU (as palmitate)/ml in multidose dispenser.	P, S, T
retinol		tablet (sugar-coated): 10 000 IU (as palmitate).	P, S, T
		water-miscible injection: 100 000 IU (as palmitate) in 2-ml ampoule.	P, S, T
riboflavin		tablet: 5 mg.	P, S, T
sodium fluoride		in any appropriate topical formulation.	P, S, T
thiamine		tablet: 50 mg (hydrochloride).	P, S, T
calcium gluconate (D)		injection: 100 mg/ml in 10-ml ampoule.	P, S, T
B complex (B1, B6 and B12)		tablets (DRAP approved)	S,T
		syrup (DRAP approved)	S,T
vitamin K1 (phytanadione)		injection: 10 mg	S,T
zinc sulfate (for acute diarrhea with ORS)		dispersible tablets: 20 mg	S,T
ferrous salt		tablets equivalent to 60 mg iron injection (sucrose salt) 20mg /ml	S,T
folic acid		syrup equivalent to 25 mg/ml iron tablets: 5mg	S,T
ferrous salt + folic acid (Nutritional supplement for use during pregnancy)		tablet, equivalent to 60 mg iron + 400 mcg folic acid	S,T
hydroxocobalamin		ampoule: 1 mg in 1ml	S,T
multiple micronutrients		sachet (UNICEF approved)	S,T
multivitamins		tablets	S,T
28. Ear, Nose, and Throat Medicines [C]			P, S, T
	acetic acid	topical: 2%, in alcohol.	

Therapeutic Category	Generic Name	Formulations	Service Level Category
<input type="checkbox"/> budenoside (D)	nasal spray: 100 micrograms per dose.	P, S, T	
<input type="checkbox"/> ciprofloxacin (D)	topical: 0.3% drops (as hydrochloride).	P, S, T	
<input type="checkbox"/> xylometazoline (a)	nasal spray: 0.05%. (a) Do not use in children less than 3 months old.	S, T	
29. Specific Medicines for Neonatal Care			
29.1. Medicines administered to the neonate [c]			
<input type="checkbox"/> caffeine citrate	injection: 20 mg/ml (equivalent to 10 mg caffeine base/ml).	S, T	
<input type="checkbox"/> chlorhexidine	oral liquid: 20 mg/ml (equivalent to 10 mg caffeine base/ml).	S, T	
	solution or gel: 7.1% (digluconate) delivering 4% chlorhexidine (for umbilical cord care) [c].	P, S, T	
<input type="checkbox"/> ibuprofen (D)	solution for injection: 5 mg/ml.	P, S, T	
<input type="checkbox"/> prostaglandin E	solution for injection: prostaglandin E1: 0.5 mg/ml in alcohol.	S, T	
	prostaglandin E 2: 1 mg/ml.	S, T	
	surfactant suspension for intratracheal instillation: 25 mg/ml or 80 mg/ml.	S, T	
29.2. Medicines administered to the mother			
<input type="checkbox"/> dexamethasone (D)	injection: 4 mg/ml dexamethasone phosphate (as disodium salt).	P, S, T	
30. Medicines for Diseases of the Joint			
30.1. Medicines used to treat gout			
<input type="checkbox"/> allopurinol	tablet: 50 mg.	S, T	
30.2. Disease-modifying agents used in rheumatoid disorders (DMARDs)			
<input type="checkbox"/> azathioprine (D)	tablet: 50 mg.	S, T	
<input type="checkbox"/> hydroxychloroquine [c]	solid oral dosage form: 200 mg (as sulfate).	S, T	
<input type="checkbox"/> methotrexate (D)	tablet: 2.5 mg (as sodium salt).	S, T	
<input type="checkbox"/> penicillamine (D)	solid oral dosage form: 250 mg.	S, T	
<input type="checkbox"/> sulfasalazine (D)	tablet: 500 mg.	S, T	
30.3. Juvenile joint diseases			
<input type="checkbox"/> acetylsalicylic acid* (acute or chronic use)	suppository: 50 mg to 150 mg. tablet: 100 mg to 500 mg. *use for rheumatic fever, juvenile arthritis, Kawasaki disease.	S, T S, T	

Medicines with Age and Weight Restriction

Table 1: Medicines with Age and Weight Restriction

atazanavir	>25 kg
atropine	>3 months
benzyl benzoate	>2 years
betamethasone topical preparations	hydrocortisone preferred in neonates
cefazolin	>1 month
ceftriaxone	>41 weeks corrected gestational age
darunavir	> 3 years
diloxanide	>25 kg
doxycycline	>8 years (except for serious infections e.g. cholera)
efavirenz	>3 years or >10 kg
emtricitabine	>3months
fluoxetine	>8 years
ibuprofen	>3 months (except IV form for patent ductus arteriosus)
mefloquine	>5 kg or >3 months
metoclopramide	Not in neonates
nevirapine	> 6 weeks
ondansetron	>1 month
saquinavir	>25 kg
silver sulfadiazine	>2 months
tetracaine	Not in preterm neonates
trimethoprim	>6 months
xylometazoline	>3 months

Index

A

- abacavir, v, 9, 26
acetazolamide, 9, 50
acetic acid, 9, 54
acetylcysteine, 9, 20
acetylsalicylic acid, 9, 18, 30, 42, 55
aciclovir, 9, 26, 49
adenosine, 9, 40
albendazole, 9, 21
alcohol-based hand rub, 9, 44
allopurinol, 9, 31, 55
all-trans retinoid acid (ATRA), v, 9, 31
aluminium hydroxide + magnesium trisilicate, 9, 46
amidotrizoate, 9, 43
amikacin, 9, 23, 25
amiloride, 9, 44
aminophylline, 9, 52
amiodarone, 9, 40
amitriptyline, 9, 18, 51
amlodipine, 9, 41
amoxicillin, 9, 21, 22
amoxicillin + clavulanic acid, 9, 22
amphotericin B, 9, 26, 29
ampicillin, 9, 10, 22
anastrozole, 9, 37
anti-D immunoglobulin, 48
anti-D immunoglobulin, 9, 39
anti-hepatitis B immunoglobulin, 9, 48
anti-rabies immunoglobulin, 48
anti-rabies immunoglobulin, 9, 39
anti-tetanus immunoglobulin, 48
anti-tetanus immunoglobulin, 9, 39
anti-venom immunoglobulin, 9, 48
artemether + lumefantrine, 9, 29
artesunate, 9, 29, 30
artesunate + sulphadoxine and pyrimethamine, 9, 30
ascorbic acid, 9, 53
asparaginase, 9, 31
atazanavir, 9, 27, 57

atazanavir + ritonavir, 27
atenolol, 9, 40, 41
atorvastatin, 9, 42
atracurium, 9, 49
atropine, 9, 17, 20, 50, 57
azathioprine, 9, 31, 55
azithromycin, 9, 23, 49
aztreonam, 9, 24

B

B complex (B1, B6 and B12, 54
barium sulfate, 9, 43
BCG vaccine, 9, 48
beclomethasone, 9, 52
bedaquiline, 9, 25
bendamustine, 9, 31
benzathine benzylpenicillin, 9, 22
benzoyl peroxide, 9, 43
benzyl benzoate, 9, 43, 57
benzylpenicillin, 9, 22
betamethasone, 9, 14, 43, 57
bevacizumab, 9, 50
bicalutamide, 9, 37
biperiden, 9, 38
bisacodyl, 9, 19, 46
bisoprolol, 9, 40, 41
bleomycin, 9, 31

Blood coagulation factors, 40
budesonide, 10, 52, 55
bupivacaine, 10, 17

C

caffeine citrate, 10, 55
calamine, 10, 43
calcium, 10, 20, 22, 31, 42, 53, 54
calcium folinate, 10, 31
calcium gluconate, 10, 20, 54

capecitabine, 10, 32
capreomycin, 10, 25
carbamazepine, 10, 20, 51
carboplatin, 10, 32
cefalexin, 10, 22
cefazolin, 10, 22, 57
cefixime, 10, 22
ceftazidime, 10, 23
ceftriaxone, 10, 22, 57
charcoal, activated, 20
chlorambucil, 10, 32
chloramphenicol, 10, 23
chlorhexidine, 10, 44, 55
chlorine base compound, 10, 44
chloroquine, 10, 30
chloroxylenol, 10, 44
chlorpheniramine, 10, 19

chlorpromazine, 10, 51
cholecalciferol, 10, 53
cholera vaccine, 10, 48
ciclosporin, 10, 31
cinnarizine, 10, 30
ciprofloxacin, 10, 23, 49, 55
cisplatin, 10, 32
clarithromycin, 10, 23
clindamycin, 10, 23
clofazimine, 10, 25
clomifene, 10, 47
clomipramine, 10, 52
clopidogrel, 10, 38, 42
clotrimazole, 10, 26
cloxacillin, 10, 22
cloxacillin + ampicillin, 22
clozapine, 10, 51
coagulation factor IX, 10, 40
coagulation factor VIII, 10, 40
coal tar, 10, 43
codeine, 10, 18
condoms, 10, 46
copper-containing device, 10, 46
cyclizine, 10, 18
cyclophosphamide, 10, 32
cycloserine, 10, 25
cytarabine, 10, 33

D

dacarbazine, 10, 33
daclatasvir, 10, 28
dactinomycin, 10, 33
darunavir, 10, 27, 57
dasabuvir, 10, 28
daunorubicin, 10, 33
deferoxamine, 10, 20, 39
delamanid, 10, 25
desmopressin, 10, 39
dexamethasone, 10, 14, 18, 19, 37, 45, 49, 50, 55
dextran 40, 10, 40
dextran 70, 10, 40
diaphragm, 10, 46
diazepam, 10, 19, 20, 51
diclofenac sodium, 10, 18
diethylcarbamazine, 10, 21
digoxin, 10, 40, 41
dihydroartemisinin + piperquine, 10, 30
diloxanide, 10, 29, 57
dimenhydrinate, 10, 46
diphtheria antitoxin, 10, 48
diphtheria vaccine, 10, 48
dobutamine, 10, 42
docetaxel, 10, 33
dolutegravir, 10, 27

dopamine, 10, 42

doxorubicin, 11, 33

doxycycline, 11, 23, 24, 30, 57

E

efavirenz (EFV or EFZ), 11, 27

enalapril, 11, 41

enoxaparin, 11, 38

entecavir, 11, 27

ephedrine, 11, 17, 42

epinephrine, 11, 12, 17, 19, 40, 50, 52

ergocalciferol, 11, 53, 54

ergometrine, 11, 50

erythromycin, 11, 49

estradiol cypionate, 11, 46

estrogen, 11, 47

ethambutol, 11, 24

ethambutol + isoniazid, 24

ethambutol + isoniazid + pyrazinamide +

rifampicin, 24

ethambutol + isoniazid + rifampicin, 24

ethanol, 11, 44

ethinylestradiol + levonorgestrel, 46

ethinylestradiol + norethisterone, 46

ethionamide, 11, 25

etonogestrel-releasing implant, 11, 46

etoposide, 11, 34

F

ferrous salt, 11, 38, 54

ferrous salt + folic acid, 11, 38, 54

filgrastim, 11, 34

fluconazole, 11, 26

flucytosine, 11, 26

fludarabine, 11, 34

fludrocortisone, 11, 46

fluorescein, 11, 43

fluorouracil, 11, 34, 43

fluoxetine, 11, 51, 57

fluphenazine, 11, 51

folic acid, 11, 38, 54

fresh-frozen plasma, 11, 39

furosemide, 11, 42, 44

G

gemcitabine, 11, 35

gentamicin, 11, 24, 49

glibenclamide, 11, 47

gliclazide, 11, 47

glucagon, 11, 47

glucose, 11, 17, 45, 53

glucose with sodium chloride, 53

glutaral, 11, 44

glycerin, 11, 46

glyceryl trinitrate, 11, 40

glycopyrrolate, 11, 17

griseofulvin, 11, 26

H

Haemophilus influenzae type b vaccine, 11,
48

haloperidol, 11, 19, 51

heparin sodium, 11, 39

hepatitis A vaccine, 11, 48

hepatitis B vaccine, 11, 48

HPV vaccine, 11, 48

hydralazine, 11, 41

hydrochlorothiazide, 11, 41, 42, 44

hydrocortisone, 11, 19, 37, 43, 45, 46, 57

hydroxocobalamin, 11, 38, 54

hydroxycarbamide, 11, 35, 39

hydroxychloroquine, 11, 55

I

ibuprofen, 11, 18, 30, 55, 57

ifosfamide, 11, 35

imatinib, 11, 35

influenza vaccine, 11, 49

insulin injection, 11, 47

intermediate-acting insulin, 11, 47

intraperitoneal dialysis solution, 51

iodine, 12, 43, 44, 47, 54

iohexol, 12, 43

ipratropium bromide, 12, 52

irinotecan, 12, 35

iron sucrose, 12, 38

isoflurane, 12, 17

isoniazid, 11, 12, 24, 27

isoniazid + pyrazinamide + rifampicin, 24

isoniazid + rifampicin, 24

isoniazid+pyridoxine+sulfamethoxazole+tr

imethoprim, 27

isoprenaline, 12, 41

isosorbide dinitrate, 12, 40

itraconazole, 12, 26

ivermectin, 12, 21

K

kanamycin, 12, 25

ketamine, 12, 17

L

labetalol, 41

lamivudine (3TC), 12, 26

latanoprost, 12, 50

ledipasvir + sofosbuvir, 28

leuprorelin, 12, 37

levamisole, 12, 21

levofloxacin, 12, 25

- levonorgestrel, 11, 12, 46, 47
levonorgestrel-releasing implant, 47
levonorgestrel-releasing intrauterine system, 46
levothyroxine, 12, 47
lidocaine, 12, 17, 40
lidocaine + epinephrine (adrenaline), 12, 17
linezolid, 12, 25
lithium carbonate, 12, 51
loperamide, 12, 19
loratadine, 12, 19
losartan potassium, 12, 40
Lugol's solution, 47
- M**
- magnesium oxides and hydroxides, 12, 46
magnesium sulfate, 12, 20
mannitol, 12, 44, 53
measles vaccine, 12, 48
mebendazole, 12, 21
medroxyprogesterone, 11, 12, 46, 47
medroxyprogesterone acetate, 11, 12, 46
mefloquine, 12, 29, 30, 57
meglumine antimoniate, 12, 29
meglumine iotroxate, 12, 44
meningococcal meningitis vaccine, 12, 48
- mercaptopurine, 12, 35
meropenem, 12, 23
mesna, 12, 35
metformin, 12, 47
methadone, 12, 52
methotrexate, 12, 35, 55
methylated spirit, 12, 44
methyldopa, 12, 41
methylprednisolone, 12, 37
methylthioninium chloride, 12, 20
metoclopramide, 12, 19, 45, 57
metronidazole, 12, 24, 29
miconazole, 12, 42, 49
midazolam, 12, 17
miltefosine, 12, 29
misoprostol, 12, 50
montelukast, 12, 52
morphine, 12, 18
moxifloxacin, 12, 23, 25
multiple micronutrients, 12, 54
multivitamins, 12, 54
mumps vaccine, 12, 49
mupirocin, 12, 42
- N**
- nalbuphine, 13, 18
naloxone, 13, 20

- naproxen, 13, 18
natamycin, 13, 49
neostigmine, 13, 49
nevirapine, 13, 27, 57
nicotinamide, 13, 54
nicotine replacement therapy (NRT), 13, 52
nifedipine, 13, 51
nitrofurantoin, 13, 24
nitrous oxide, 13, 17
norethisterone enantate, 13, 46
normal immunoglobulin, 13, 39
nystatin, 13, 26
- O**
- ofloxacin, 13, 25, 49
olanzapine, 13, 51
omeprazole, 13, 44
ondansetron, 13, 19, 45, 57
oral rehydration salts, 13, 45, 53
oxaliplatin, 13, 36
oxygen, 13, 17
oxytocin, 13, 50, 51
- P**
- paclitaxel, 13, 36
p-aminosalicylic acid, 13, 25
- pancreatic enzymes, 13, 44
pancuronium, 13, 49
paracetamol, 13, 18, 30
paromomycin, 13, 29
pegylated interferon alfa (2a or 2b), 13, 28
penicillamine, 13, 55
permethrin, 13, 43
pertussis vaccine, 13, 48
pheniramine, 13, 20
phenobarbital, 13, 20
phenoxymethylenicillin, 13, 22
phenytoin, 13, 21
phloroglucinol, 13, 46
phytomenadione, 13, 39
pilocarpine, 13, 50
piperacillin + tazobactum, 13, 23
platelets, 13, 39
pneumococcal vaccine, 13, 48
podophyllum resin, 13, 43
poliomyelitis vaccine, 13, 48
polymyxin B (sulphate)+ bacitracin zinc, 13, 50
potassium chloride, 13, 53
potassium iodide, 13, 47
potassium permanganate, 13, 42
povidone iodine, 13, 44
praziquantel, 13, 21

prednisolone, 13, 19, 38, 50
pregabalin, 13, 19
primaquine, 13, 29
procaine benzyl penicillin, 13, 22
procarbazine, 13, 36
progesterone vaginal ring, 13, 47
proguanil, 13, 30
proparacaine, 13, 50
propofol, 13, 17
propranolol, 13, 30, 40, 41
propylthiouracil, 13, 47
prostaglandin E, 13, 55
protamine sulfate, 13, 39
pyrantel, 13, 21
pyrazinamide, 11, 12, 13, 24, 25
pyridostigmine, 13, 49
pyridoxine, 12, 13, 27, 54

R

rabies vaccine, 13, 49
raltegravir, 13, 27
ranitidine, 13, 44
red blood cells, 13, 39
retinol, 13, 54
ribavirin, 13, 27, 28
riboflavin, 14, 54
rifabutin, 14, 25

rifampicin, 11, 12, 14, 25
rifapentine, 14, 25
risovastatin, 14, 42
risperidone, 14, 51
ritonavir, 9, 13, 14, 27, 28
rituximab, 14, 36
rotavirus vaccine, 14, 48
rubella vaccine, 14, 48

S

salbutamol, 14, 52
salicylic acid, 14, 43
salicylic acid + betamethasone, 43
selenium sulfide, 14, 42
senna, 14, 45
sevoflurane, 14, 17
silver sulfadiazine, 14, 42, 43, 57
simeprevir, 14, 28
simvastatin, 14, 42
sodium biphosphate, 14, 46
sodium chloride, 11, 14, 45, 53
sodium cromoglycate, 14, 49
sodium fluoride, 14, 54
sodium hydrogen carbonate, 14, 53
sodium lactate, compound solution, 53
sodium nitrite, 14, 20
sodium nitroprusside, 14, 41

- sodium polystyrene sulfonate, 14, 44
sodium stibogluconate, 14, 29
sodium thiosulfate, 14, 42, 43
sofosbuvir, 12, 14, 28
spectinomycin, 14, 24
spironolactone, 14, 42, 44
streptokinase, 14, 42
streptomycin, 14, 25
sulfadoxine + pyrimethamine, 14, 29, 30
sulfamethoxazole + trimethoprim, 14, 24
sulfasalazine, 14, 45, 55
sumatriptan, 14, 30
surfactant, 14, 55
suxamethonium, 14, 49
- thioguanine, 14, 36
timolol, 14, 50
tirofiban, 14, 38
tobramycin + dexamethasone, 50
tramadol, 14, 19
tranexamic acid, 14, 39
trastuzumab, 14, 36
tropicamide, 14, 43, 50
tuberculin, purified protein derivative (PPD), 47
typhoid vaccine, 14, 49

T

- tamoxifen, 14, 38
tears natural, 14, 50
tenofovir disoproxil fumarate (TDF), 14, 28
terbinafine, 14, 42
testosterone, 14, 46
tetanus immunoglobulin, 14, 48
tetanus vaccine, 14, 48
tetracaine, 14, 50, 57
tetracycline, 14, 49
thiamine, 14, 54

U

- urea, 14, 43

V

- valproic acid, 14, 21, 51
vancomycin, 14, 24
varicella vaccine, 14, 49
vecuronium, 14, 49
verapamil, 15, 40
vinblastine, 15, 36
vincristine, 15, 36
vinorelbine, 15, 37
vitamin K1 (phytonadione), 15, 54

W

- warfarin, 15, 39
- water for injection, 15, 53
- whole blood, 15, 39

X

- xylometazoline, 15, 55, 57

Y

- yellow fever vaccine, 15, 48

Z

- zidovudine (ZDV or AZT)., 15, 27
- zinc sulfate, 15, 45, 54

Application Procedure for Inclusion, Deletion or Modification of a Medicine in the Next Edition of National Essential Medicine List, 2020

Applications, recommendation and suggestions for inclusion, deletion, change i.e addition or modification of an indication of a medicine in the next edition of National Essential Medicine List (NEML, 2020) should be forwarded to the Division of Pharmacy Services, Drug Regulatory Authority of Pakistan, Islamabad, in a format as prescribed below;

Information to be included with an application for inclusion, deletion or modification of a medicine in the National Essential Medicine List (NEML).

1. Summary statement of the proposal for inclusion, change or deletion
2. Name of the technical expert(s) consulted (if applicable)
3. Name of the organization(s) consulted and/or supporting the application
4. International Nonproprietary Name (INN, generic name) of the medicine
5. Formulation proposed for inclusion; including adult and paediatric (if appropriate)
6. International and local availability including name of manufacturers and trade names
7. Whether listing is requested as an individual medicine or as an example of a therapeutic group
8. Information supporting the public health relevance (epidemiological information on disease burden, assessment of current use, target population)
9. Treatment details (dosage regimen, duration; reference to existing WHO and other clinical guidelines; need for special diagnostics, treatment or monitoring facilities and skills)

10. Summary of comparative effectiveness in a variety of clinical settings:
 - Identification of clinical evidence (search strategy, systematic reviews identified, reasons for selection/exclusion of particular data)
 - Summary of available data (appraisal of quality, outcome measures, summary of results)
 - Summary of available estimates of comparative effectiveness
11. Summary of comparative evidence on safety:
 - Estimate of total patient exposure to date
 - Description of the adverse effects/reactions and estimates of their frequency
 - Identification of variation in safety that may relate to health systems and patient factors
 - Summary of comparative safety against comparators
12. Summary of available data on comparative cost and cost-effectiveness within the pharmacological class or therapeutic group:
 - range of costs of the proposed medicine
 - resource use and comparative cost-effectiveness presented as range of cost per routine outcome
13. Summary of regulatory status of the medicine (in various countries)
14. Availability of pharmacopoeial standards (British Pharmacopoeia, International Pharmacopoeia, United States Pharmacopoeia, European Pharmacopeia)
15. Proposed (new/adapted) text that could be included in the revised version.

For further information, please contact:

Pharmacy Services Division,
Drug Regulatory Authority of Pakistan,
T.F Complex,
7 Mauve Area, Sector G-9/4, Islamabad
Phone No. 051-9262160
Email: dir.ps@dra.gov.pk
Website: www.dra.gov.pk

References

1. National Essential Medicines List – 2016, Drug Regulatory Authority of Pakistan, Ministry of National Health Services, Regulations and Coordination, Government of Pakistan
2. World Health Organization (WHO) Model List of Medicines (20th list), March 2017
3. Ministry of National Health Services, Regulations and Coordination [2014-18] Comprehensive Multiyear Plan; Immunization Programme in Pakistan. (2014-2018). [Islamabad]
4. Directorate of Malaria; Ministry of Health (MOH), Pakistan. 2007. Malaria Case Management; Desk guide for clinician and Health care providers. Islamabad: MOH.
5. National TB Control Program, Islamabad. (Jan 2015). National Guidelines for the control of Tuberculosis in Pakistan.
6. World Health Organization country office; Pakistan. (2014). Pakistan Pharmaceutical Situation Assessment Level II. Islamabad



USAID
FROM THE AMERICAN PEOPLE

**USAID Global Health Supply Chain Program
Procurement and Supply Management**



**World Health
Organization**

